


What is claimed is:

1. A compound of formula (I):



wherein:

q is an integer from 1 to 4;

n is an integer from 0 to 2;

R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituents selected from the group consisting of OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, N<sup>+</sup>R<sup>9</sup>R<sup>10</sup>R<sup>w</sup>A<sup>-</sup>, SR<sup>9</sup>, S<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, P<sup>+</sup>R<sup>9</sup>R<sup>10</sup>R<sup>11</sup>A<sup>-</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, SO<sub>3</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, CN, halogen, oxo, and CONR<sup>9</sup>R<sup>10</sup>,

wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl optionally have one or more carbons replaced by O, NR<sup>9</sup>,

$N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^W$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, arylalkyl, and alkylammoniumalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which they are attached form C<sub>3</sub>-C<sub>10</sub> cycloalkyl;

$R^3$  and  $R^4$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, and SO<sub>3</sub>R<sup>9</sup>, wherein  $R^9$  and  $R^{10}$  are as defined above; or

$R^3$  and  $R^4$  together form =O, =NOR<sup>11</sup>, =S, =NNR<sup>11</sup>R<sup>12</sup>, =NR<sup>9</sup>, or =CR<sup>11</sup>R<sup>12</sup>,

wherein  $R^{11}$  and  $R^{12}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, SO<sub>3</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, CN, halogen, oxo, and CONR<sup>9</sup>R<sup>10</sup>, wherein  $R^9$  and  $R^{10}$  are as defined above, provided that both  $R^3$  and  $R^4$  cannot be OH, NH<sub>2</sub>, and SH, or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

$R^5$  and  $R^6$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, OR<sup>30</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, and SO<sub>3</sub>R<sup>9</sup>,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl,

heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,

wherein:

$A^-$  is a pharmaceutically acceptable anion and  $M$  is a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,  $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  $P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, aryl,

arylalkyl, cycloalkyl, heterocycle, heteroaryl,  
quaternary heterocycle, quaternary heteroaryl, and  
quaternary heteroarylalkyl,

5 wherein alkyl, alkenyl, alkynyl, arylalkyl,  
heterocycle, and polyalkyl optionally have one or more  
carbons replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  
 $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ ,  $\text{P}(\text{O})\text{R}^9$ , phenylene, carbohydrate,  
amino acid, peptide, or polypeptide, and

10  $\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are optionally substituted with  
one or more groups selected from the group consisting  
of sulfoalkyl, quaternary heterocycle, quaternary  
heteroaryl,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
 $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ , CN, halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_2\text{OM}$ ,  
15  $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{R}^{11}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , and  
 $\text{C}(\text{O})\text{OM}$ ,

wherein  $\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected  
from the substituents constituting  $\text{R}^9$  and M; or

$\text{R}^{14}$  and  $\text{R}^{15}$ , together with the nitrogen atom to  
which they are attached, form a cyclic ring; and  
20  $\text{R}^{30}$  is selected from the group consisting of alkyl,  
alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; and

$\text{R}^7$  and  $\text{R}^8$  are independently selected from the  
group consisting of hydrogen and alkyl; and

25 one or more  $\text{R}^x$  are independently selected from the  
group consisting of H, alkyl, alkenyl, alkynyl,  
polyalkyl, acyloxy, aryl, arylalkyl, halogen,  
haloalkyl, cycloalkyl, heterocycle, heteroaryl,  
polyether, quaternary heterocycle, quaternary  
30 heteroaryl,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S}(\text{O})\text{R}^{13}$ ,  $\text{S}(\text{O})_2\text{R}^{13}$ ,



5  $\text{SO}_3\text{R}^{13}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ ,  
 $\text{CN}$ ,  $\text{OM}$ ,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$ ,  $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  
 $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$ ,  $\text{C}(\text{O})\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{OR}^{18}$ ,  $\text{S}(\text{O})_n\text{NR}^{18}$ ,  $\text{NR}^{13}\text{R}^{18}$ ,  
 $\text{NR}^{18}\text{OR}^{14}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{P}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ , amino acid,  
peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl,  
polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl,  
polyether, quaternary heterocycle, and quaternary  
heteroaryl can be further substituted with  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  
10  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ ,  
 $\text{CN}$ , halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ ,  
 $\text{P}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{C}(\text{O})\text{OM}$ , and

wherein  $\text{R}^{18}$  is selected from the group consisting  
of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle,  
15 heteroaryl, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl,  
heterocycle, heteroaryl, alkyl, quaternary heterocycle,  
and quaternary heteroaryl optionally are substituted  
with one or more substituents selected from the group  
20 consisting of  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
 $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ ,  $\text{CN}$ , halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_3\text{R}^9$ ,  
 $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ , and  $\text{C}(\text{O})\text{OM}$ ,

wherein in  $\text{R}^x$ , one or more carbons are optionally  
replaced by O,  $\text{NR}^{13}$ ,  $\text{N}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^{13}\text{A}^-$ ,  
25  $\text{PR}^{13}$ ,  $\text{P}(\text{O})\text{R}^{13}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , phenylene, amino acid,  
peptide, polypeptide, carbohydrate, polyether, or  
polyalkyl,

wherein in said polyalkyl, phenylene, amino acid,  
peptide, polypeptide, and carbohydrate, one or more

carbons are optionally replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ ,  
S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{P}(\text{O})\text{R}^9$ ;

wherein quaternary heterocycle and quaternary  
heteroaryl are optionally substituted with one or more  
5 groups selected from the group consisting of alkyl,  
alkenyl, alkynyl, polyalkyl, polyether, aryl,  
haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,  
oxo,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S}(\text{O})\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  
 $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ , CN, OM,  $\text{SO}_2\text{OM}$ ,  
10  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{P}(\text{O})\text{R}^{13}\text{R}^{14}$ ,  
 $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  $\text{P}(\text{OR}^{13})\text{OR}^{14}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  
 $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,

provided that both  $\text{R}^5$  and  $\text{R}^6$  cannot be hydrogen or  
SH;

15 provided that when  $\text{R}^5$  or  $\text{R}^6$  is phenyl, only one of  
 $\text{R}^1$  or  $\text{R}^2$  is H;

provided that when  $q = 1$  and  $\text{R}^x$  is styryl,  
anilido, or anilinocarbonyl, only one of  $\text{R}^5$  or  $\text{R}^6$  is  
alkyl; or

20 a pharmaceutically acceptable salt, solvate, or  
prodrug thereof.

2. A compound of claim 1, wherein  $\text{R}^5$  and  $\text{R}^6$  are  
25 independently selected from the group consisting of H,  
aryl, heterocycle, quaternary heterocycle, and  
quaternary heteroaryl,


wherein said aryl, heteroaryl, quaternary  
heterocycle, and quaternary heteroaryl can be  
30 substituted with one or more substituent groups

independently selected from the group consisting of  
alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl,  
haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,  
oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  
5  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  
 $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  
 $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  
 $N^+R^9R^{11}R^{12}A^-$ ,

wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
10 polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can optionally have one or more carbons replaced by O,  
 $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  
 $P^+R^7R^8A^-$ , or phenylene,

wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
15 polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can be further substituted with one or more substituent  
groups selected from the group consisting of  $OR^7$ ,  
 $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  
 $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl,  
20 cycloalkyl, heterocycle, arylalkyl, quaternary  
heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8A^-$ ,  
and  $P(O)(OR^7)OR^8$ .

3. A compound of claim 2, wherein  $R^5$  or  $R^6$  has  
25 the formula



wherein:

t is an integer from 0 to 5;

Ar is selected from the group consisting of phenyl, thiophenyl, pyridyl, piperazinyl, piperonyl, pyrrolyl, naphthyl, furanyl, anthracenyl, quinolinyl, isoquinolinyl, quinoxalinyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, pyrimidinyl, thiazolyl, triazolyl, isothiazolyl, indolyl, benzoimidazolyl, benzoxazolyl, benzothiazolyl, and benzoisothiazolyl; and

one or more  $R^Y$  are independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,


wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,  $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,

$P^+R^7R^8A^-$ , or phenylene.

4. A compound of claim 3, wherein  $R^5$  or  $R^6$  has the formula (II)

5



(II)

5. A compound of claim 4, wherein  $n$  is 1 or 2.

10

6. A compound of claim 5, wherein one of  $R^7$  or  $R^8$  is H and the other of  $R^7$  or  $R^8$  is alkyl.

7. A compound of claim 5, wherein both  $R^7$  and  $R^8$  are H.

15

8. A compound of claim 7, wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of H and alkyl.

20

9. A compound of claim 8, wherein said alkyl is a  $C_1$ - $C_{10}$  alkyl.

10. A compound of claim 8, wherein  $R^1$  and  $R^2$  are both alkyl.

25

11. A compound of claim 10, wherein said alkyl is a  $C_1$ - $C_{10}$  alkyl.

12. A compound of claim 11, wherein said alkyl is  
a C<sub>2</sub>-C<sub>7</sub> alkyl.

5 13. A compound of claim 12, wherein said alkyl is  
a C<sub>2</sub>-C<sub>4</sub> alkyl.

10 14. A compound of claim 13, wherein said alkyl is  
independently selected from the group consisting of  
ethyl, n-propyl, n-butyl, and isobutyl.

15 15. A compound of claim 8, wherein R<sup>1</sup> and R<sup>2</sup> are  
each n-butyl.

16. A compound of claim 8, wherein one of R<sup>1</sup> and  
R<sup>2</sup> is ethyl and the other of R<sup>1</sup> and R<sup>2</sup> is n-butyl.

17. A compound of claim 15, wherein q is 1, 2, or  
3.

18. A compound of claim 16, wherein q is 1, 2, or  
3.

19. A compound of claim 17, wherein q is 1 or 2.

20. A compound of claim 19, wherein q is 1.

21. A compound of claim 18, wherein q is 1 or 2.

22. A compound of claim 21, wherein q is 1.

23. A compound of claim 19, wherein R<sup>3</sup> and R<sup>4</sup> are  
independently selected from the group consisting of H

and OR<sup>9</sup>.

24. A compound of claim 21, wherein R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of H and OR<sup>9</sup>.

25. A compound of claim 23, wherein R<sup>9</sup> is H.

26. A compound of claim 24, wherein R<sup>9</sup> is H.

27. A compound of claim 25, wherein one or more R<sup>x</sup> are in the 7-, 8-, or 9-position of the benzo ring of formula (I).

28. A compound of claim 26, wherein said R<sup>x</sup> is in the 7-, 8-, or 9- position of the benzo ring of formula (I).

29. A compound of claim 27, wherein said R<sup>x</sup> are in the 7- and 9- positions of the benzo ring of formula (I).

30. A compound of claim 28, wherein said R<sup>x</sup> is in the 7-position of the benzo ring of formula (I).

31. A compound of claim 29, wherein said one or more R<sup>x</sup> are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, alcyloxy, polyether, halogen, OR<sup>13</sup>, NR<sup>13</sup>R<sup>14</sup>, NR<sup>13</sup>NR<sup>14</sup>R<sup>15</sup>, N<sup>+</sup>R<sup>9</sup>R<sup>11</sup>R<sup>12</sup>A<sup>-</sup>, SR<sup>13</sup>, S<sup>+</sup>R<sup>13</sup>R<sup>14</sup>, CO<sub>2</sub>R<sup>13</sup>, NR<sup>14</sup>C(O)R<sup>13</sup>, and NR<sup>14</sup>C(O)R<sup>13</sup>,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ .

32. A compound of claim 30, wherein said  $R^X$  is selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^{13}$ ,  $S^+R^{13}R^{14}$ ,  $CO_2R^{13}$ ,  $NR^{14}C(O)R^{13}$ , and  $NR^{14}C(O)R^{13}$ ,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,



$PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

5 wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ .

10 33. A compound of claim 31, wherein said one or more Rx are independently selected from the group consisting of polyether,  $OR^{13}$ ,  $NR^{13}R^{14}$ , and  $N^+R^9R^{11}R^{12}A^-$ .

15 34. A compound of the claim 32, wherein said  $R^x$  is selected from the group consisting of polyether,  $OR^{13}$ ,  $NR^{13}R^{14}$ , and  $N^+R^9R^{11}R^{12}A^-$ .

20 35. A compound of claim 33, wherein said one or more Rx are independently selected from the group consisting of  $OR^{13}$  and  $NR^{13}R^{14}$ .

25 36. A compound of claim 34, wherein said  $R^x$  is independently selected from the group consisting of  $OR^{13}$  and  $NR^{13}R^{14}$ .

37. A compound of claim 35, wherein  $R^{13}$  and  $R^{14}$  each methyl.

30 38. A compound of the claim 36, wherein  $R^{13}$  and  $R^{14}$  each methyl.

39. A compound of claim 31, wherein one or more  $R^Y$  are independently in the 3- or the 4-position of the phenyl ring of formula (II).

40. A compound of claim 32, wherein one or more  $R^Y$  are independently in the 3- or the 4- position of the phenyl ring of formula (II).

41. A compound of claim 39, wherein t is 1 or 2.

42. A compound of claim 40, wherein t is 1 or 2.

43. A compound of claim 41, wherein said one or more  $R^Y$  are independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide,  $NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ , and  $OR^{13}$ , wherein alkyl and polyether can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

44. A compound of claim 42, wherein said  $R^Y$  is independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide,  $NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ , and  $OR^{13}$ , wherein alkyl and polyether can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

45. A compound of claim 43, wherein said one or

more  $R^Y$  are independently selected from the group consisting of alkyl, polyether, fluoride,  $NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ , and  $OR^{13}$ ,

5 wherein alkyl and polyether can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

10 46. A compound of claim 44 wherein said  $R^Y$  is independently selected from the group consisting of alkyl, polyether, fluoride,  $NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ , and  $OR^{13}$ ,

15 wherein alkyl and polyether can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

47. A compound of claim 45, wherein said  $R^{13}$  and  $R^{14}$  are alkyl,

20 wherein alkyl can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

48. A compound of claim 46, wherein said  $R^9$  and  $R^{10}$  are alkyl,

25 wherein alkyl can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

49. A compound of claim 47, wherein  $n$  is 2.


50. A compound of claim 48, wherein  $n$  is 2.

30 51. A compound of claim 49, wherein said OH group is in a syn relationship to said structure of formula (II).

52. A compound of claim 50, wherein said OH group is in a *syn* relationship to said structure of formula (II).


5

53. A compound of claim 51, having the formula:



10


54. A compound of claim 51, having the formula:



SRL 6071  
(C-3021/2)


540

55. A compound of claim 51, having the formula:



5


56. A compound of claim 51, having the formula:



SRL 6071  
(C-3021/2)


541

57. A compound of claim 51, having the formula:



5


58. A compound of claim 52, having the formula:



SRL 6071  
(C-3021/2)


542

59. A compound of claim 52, having the formula:




5

60. A compound of claim 52, having the formula:




61. A compound of claim 52, having the formula:



5

62. A compound of claim 52, having the formula:




10

63. A compound of claim 31, wherein  $n$  is 1.

64. A compound of claim 63, wherein  $R^y$  is H.

65. A compound of claim 64, having the formula





5 66. A compound of claim 4, wherein R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of H and alkyl.

67. A compound of claim 66, wherein said alkyl is C<sub>1</sub>-C<sub>10</sub> alkyl.

10 68. A compound of claim 67, wherein said alkyl is C<sub>2</sub>-C<sub>7</sub> alkyl.

69. A compound of claim 68, wherein said alkyl is C<sub>2</sub>-C<sub>4</sub> alkyl.

15 70. A compound of claim 69, wherein R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.

20 71. A compound of claim 4, wherein R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of H and OR<sup>9</sup>.

72. A compound of claim 71, wherein R<sup>9</sup> is H.

73. A compound of claim 4, wherein n is 2.

5 74. A compound of claim 3, wherein  $R^3$  and  $R^4$  are independently selected from the group consisting of H and  $OR^9$ .

75. A compound of claim 74, wherein  $R^9$  is H.

10 76. A compound of claim 3, wherein one of  $R^7$  or  $R^8$  is H.

15 77. A compound of claim 76, wherein both  $R^7$  and  $R^8$  are H.

78. A compound of claim 3, wherein said one or more  $R^x$  are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^{13}$ ,  $S^+R^{13}R^{14}$ ,  $CO_2R^{13}$ ,  $NR^{14}C(O)R^{13}$ , and  $NR^{14}C(O)R^{13}$ ,

20 wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

25 wherein in  $R^x$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid,

30

peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more  
5 carbons are optionally replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{P}(\text{O})\text{R}^9$ .

79. A compound of claim 78, wherein said one or more  $\text{R}^x$  are independently selected from the group  
10 consisting of polyether,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ .

80. A compound of claim 79, wherein said one or more  $\text{R}^x$  are independently selected from the group  
15 consisting of  $\text{OR}^{13}$  and  $\text{NR}^{13}\text{R}^{14}$ .

81. A compound of claim 80, wherein  $\text{R}^{13}$  and  $\text{R}^{14}$  are each methyl.

82. A compound of claim 3, wherein one or more  $\text{R}^y$  are independently in the 3- or the 4-position of the phenyl ring of formula (II).

83. A compound of claim 82, wherein one or more  $\text{R}^y$  is selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide,  $\text{NR}^9\text{R}^{10}$ , and  $\text{NC}(\text{O})\text{R}^9$ ,

wherein alkyl and polyether can be substituted with  $\text{SO}_3\text{R}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ , and quaternary heteroaryl.

30

84. A compound of claim 83, wherein  $\text{R}^9$  and  $\text{R}^{10}$

are alkyl.

85. A compound of claim 84, wherein one or more  $R^Y$  is selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide,  $NR^9R^{10}$ , and  $NC(O)R^9$ .

86. A compound of claim 1, wherein said one or more  $R^X$  are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^{13}$ ,  $S^+R^{13}R^{14}$ ,  $CO_2R^{13}$ ,  $NR^{14}C(O)R^{13}$ , and  $NR^{14}C(O)R^{13}$ ,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ .

87. A compound of claim 1, wherein n is 1 or 2.

88. A compound of claim 87, wherein n is 2.)

5 89. A compound of claim 1, wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of H and alkyl.

10 90. A compound of claim 89, wherein said alkyl is  $C_1$ - $C_{10}$  alkyl.

91. A compound of claim 90, wherein said alkyl is  $C_2$ - $C_7$  alkyl.

15 92. A compound of claim 91, wherein said alkyl is  $C_2$ - $C_4$  alkyl.

20 93. A compound of claim 92, wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.


94. A compound of claim 1, wherein  $R^3$  and  $R^4$  are independently selected from the group consisting of H and  $OR^9$ .

25 95. A compound of claim 94, wherein  $R^9$  is H.

96. A compound of claim 1, wherein one of  $R^7$  or  $R^8$  is H.

30 97. A compound of claim 96, wherein both  $R^7$  and  $R^8$  are H.

98. A compound of the formula (III)



wherein :

q and r are independently integers from 0 to 4;

d and e are independently integers from 0 to 2;

t and u are independently integers from 0 to 4;

$R^1$ ,  $R^{1A}$ ,  $R^2$ , and  $R^{2A}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituent selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^WA^-$ ,  $SR^9$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, polyalkyl, aryl, and cycloalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^W$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which they are attached form  $C_3$ - $C_{10}$  cycloalkylidene, or

$R^{1A}$  and  $R^{2A}$  taken together with the carbon to which they are attached form  $C_3$ - $C_{10}$  cycloalkylidene;

5  $R^3$ ,  $R^{3A}$ ,  $R^4$ , and  $R^{4A}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ , wherein  $R^9$  and  $R^{10}$  are as defined above; or

10  $R^3$  and  $R^4$  together form  $=O$ ,  $=NOR^{11}$ ,  $=S$ ,  $=NNR^{11}R^{12}$ ,  $=NR^9$ , or  $=CR^{11}R^{12}$ , or

$R^{3A}$  and  $R^{4A}$  together form  $=O$ ,  $=NOR^{11}$ ,  $=S$ ,  $=NNR^{11}R^{12}$ ,  $=NR^9$ , or  $=CR^{11}R^{12}$ ,

15 wherein  $R^{11}$  and  $R^{12}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carbalkoxyalkyl, cycloalkyl, cyanoalkyl,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,  
20 wherein  $R^9$  and  $R^{10}$  are as defined above, provided that both  $R^3$  and  $R^4$  cannot be OH, NH<sub>2</sub>, and SH, or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

25 wherein  $A^-$  is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation;

$R^7$ ,  $R^{7A}$ ,  $R^8$ , and  $R^{8A}$  are independently selected from the group consisting of hydrogen and alkyl; and

30 one or more  $R^X$  and  $R^{XA}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen,

haloalkyl, cycloalkyl, heterocycle, heterocycle,  
polyether, quaternary heterocycle, quaternary  
heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  
 $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  
5  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  
 $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  
 $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid,  
peptide, polypeptide, and carbohydrate,

10 wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl,  
polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl,  
polyether, quaternary heterocycle, and quaternary  
heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  
 $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  
 $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  
15  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting  
of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle,  
heterocycle, alkyl,

20 wherein acyl, arylalkoxycarbonyl, arylalkyl,  
heterocycle, heterocycle, alkyl quaternary heterocycle,  
and quaternary heteroaryl optionally are substituted  
with one or more substituent selected from the group  
consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  
 $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  
25  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

wherein in  $R^*$  and  $R^{xa}$ , one or more carbons are  
optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  
 $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino  
acid, peptide, polypeptide, carbohydrate, polyether, or



polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ ,  
5 S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{P}(\text{O})\text{R}^9$ ;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl,  
10 haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S}(\text{O})\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ , CN, OM,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{P}(\text{O})\text{R}^{13}\text{R}^{14}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  $\text{P}(\text{OR}^{13})\text{OR}^{14}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,

15  $\text{R}^{19}$  is selected from the group consisting of alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide, polypeptide, wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate,  
20 amino acid, and peptide, polypeptide, can optionally have one or more carbon replaced by O,  $\text{NR}^7$ ,  $\text{N}^+\text{R}^7\text{R}^8$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^7\text{R}^8$ ,  $\text{PR}^7$ ,  $\text{P}^+\text{R}^7\text{R}^8$ , phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl,

25 wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, peptide, and polypeptide can be substituted with one or more substituent groups independently selected from the  
30 group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle,

arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  
 $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  
 $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  
 $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  
5  $N^+R^9R^{11}R^{12}A^-$ ;

wherein one or more  $R^y$  and  $R^{yA}$  are independently  
selected from the group consisting of H, alkyl,  
alkenyl, alkynyl, aryl, cycloalkyl, heterocycle,  
quaternary heterocycle,  $OR^9$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  
10  $SO_3R^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl,  
and heterocycle can be substituted with one or more  
substituent groups independently selected from the  
group consisting of alkyl, alkenyl, alkynyl, polyalkyl,  
15 polyether, aryl, haloalkyl, cycloalkyl, heterocycle,  
arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  
 $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  
 $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  
 $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  
20  $N^+R^9R^{11}R^{12}A^-$ ,

wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can be further substituted with one or more substituent  
groups selected from the group consisting of  $OR^7$ ,  
25  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  
 $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl,  
cycloalkyl, heterocycle, arylalkyl, quaternary  
heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8A^-$ ,  
and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can optionally have one or more carbons replaced by O,  
NR<sup>7</sup>, N<sup>+</sup>R<sup>7</sup>R<sup>8</sup>A-, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>7</sup>A-, PR<sup>7</sup>, P(O)R<sup>7</sup>,  
5 P<sup>+</sup>R<sup>7</sup>R<sup>8</sup>A-, or phenylene.

99. A compound of claim 98, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently selected from the group  
consisting of H and alkyl.

100. A compound of claim 99, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently selected from the group  
consisting of H and C<sub>1</sub>-C<sub>10</sub> alkyl.

101. A compound of claim 100, wherein said alkyl  
is a C<sub>2</sub>-C<sub>7</sub> alkyl.

102. A compound of claim 101, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently C<sub>2</sub>-C<sub>4</sub> alkyl.

103. A compound of claim 102, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently selected from the group  
consisting of ethyl, n-propyl, n-butyl, and isobutyl.

104. A compound of claim 98, wherein R<sup>3</sup>, R<sup>3A</sup>, R<sup>4</sup>,  
and R<sup>4A</sup> are independently selected from the group  
consisting of H and OR<sup>9</sup>.

105. A compound of claim 104, wherein R<sup>9</sup> is H.

106. A compound of claim 98, wherein R<sup>7</sup>, R<sup>7A</sup>, R<sup>8</sup>,  
and R<sup>8A</sup> are H.

107. A compound of claim 98, wherein d and e are

independently 1 or 2.

108. A compound of claim 107, wherein d and e are both 2.

5

109. A compound of claim 98, wherein one or more  $R^X$  and one or more  $R^{xA}$  are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^{13}$ ,  $S^+R^{13}R^{14}$ ,  $CO_2R^{13}$ ,  $NR^{14}C(O)R^{13}$ , and  $NR^{14}C(O)R^{13}$ ,

10

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

15

wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

20

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ .

25

110. A compound of claim 98, wherein one or more  $R^Y$  and one or more  $R^{yA}$  are independently selected from the group consisting of alkyl, polyether, fluoride,

30

chloride, bromide, iodide,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$ , and  $\text{OR}^{13}$ ,

wherein alkyl and polyether can be further substituted with  $\text{SO}_3\text{R}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ , and quaternary heteroaryl.

111. A compound of claim 98, wherein  $\text{R}^{19}$  is selected from the group consisting of alkane diyl, polyalkane diyl, alkoxy diyl, and polyalkoxy diyl, wherein alkane diyl and polyalkane diyl can optionally have one or more carbon replaced by O,  $\text{NR}^7$ ,  $\text{N}^+\text{R}^7\text{R}^8$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^7\text{R}^8$ ,  $\text{PR}^7$ ,  $\text{P}^+\text{R}^7\text{R}^8$ , or phenylene.

112. A compound of claim 111, wherein  $\text{R}^{19}$  is selected from the group consisting of alkoxy diyl and polyalkoxydiyl wherein one or more carbons are optionally replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{R}^{10}$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, or polyalkyl.

113. A compound of claim 112, wherein  $\text{R}^1$ ,  $\text{R}^{1A}$ ,  $\text{R}^2$ , and  $\text{R}^{2A}$  are independently selected from the group consisting of H and alkyl.

114. A compound of claim 113, wherein  $\text{R}^3$ ,  $\text{R}^{3A}$ ,  $\text{R}^4$ , and  $\text{R}^{4A}$  are independently selected from the group consisting of H and  $\text{OR}^9$ .

115. A compound of claim 114, wherein  $\text{R}^9$  is H.

116. A compound of claim 115, wherein  $\text{R}^7$ ,  $\text{R}^{7A}$ ,  $\text{R}^8$ , and  $\text{R}^{8A}$  are each H.

117. A compound of claim 116, wherein d and e are independently 1 or 2.

118. A compound of claim 117, wherein one or more  
5  $R^X$  and one or more  $R^{XA}$  are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^{13}$ ,  $S^+R^{13}R^{14}$ ,  $CO_2R^{13}$ ,  $NR^{14}C(O)R^{13}$ , and  $NR^{14}C(O)R^{13}$ ,

10 wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  
15  $C(O)OM$ , and


wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or  
20 polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ .

119. A compound of claim 118, wherein one or more  
25  $R^Y$  and one or more  $R^{YA}$  are independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide,  $NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ , and  
30  $OR^{13}$ ,


wherein alkyl and polyether can be further substituted with  $\text{SO}_3\text{R}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ , and quaternary heteroaryl.

5 120. A compound of claim 119, having the formula:



PEG = 3400 molecular weight polyethylene glycol polymer chain

121. A compound of the formula (IV)



5 wherein :

q and r are independently integers from 0 to 3;

d and e are independently integers from 0 to 2;

t and u are independently integers from 0 to 5;

10  $R^1$ ,  $R^{1A}$ ,  $R^2$ , and  $R^{2A}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

15 wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituent selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^{11}A^-$ ,  $SR^9$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

20 wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, polyalkyl, aryl, and cycloalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,



$N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^W$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which they are attached form C<sub>3</sub>-C<sub>10</sub> cycloalkylidene, or

$R^{1A}$  and  $R^{2A}$  taken together with the carbon to which they are attached form C<sub>3</sub>-C<sub>10</sub> cycloalkylidene;

$R^3$ ,  $R^{3A}$ ,  $R^4$ , and  $R^{4A}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, and SO<sub>3</sub>R<sup>9</sup>, wherein R<sup>9</sup> and R<sup>10</sup> are as defined above; or

$R^3$  and  $R^4$  together form =O, =NOR<sup>11</sup>, =S, =NNR<sup>11</sup>R<sup>12</sup>, =NR<sup>9</sup>, or =CR<sup>11</sup>R<sup>12</sup>, or

$R^{3A}$  and  $R^{4A}$  together form =O, =NOR<sup>11</sup>, =S, =NNR<sup>11</sup>R<sup>12</sup>, =NR<sup>9</sup>, or =CR<sup>11</sup>R<sup>12</sup>,

wherein  $R^{11}$  and  $R^{12}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl,

cycloalkyl, cyanoalkyl, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, SO<sub>3</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, CN, halogen, oxo, and CONR<sup>9</sup>R<sup>10</sup>,

wherein R<sup>9</sup> and R<sup>10</sup> are as defined above, provided that both R<sup>3</sup> and R<sup>4</sup> cannot be OH, NH<sub>2</sub>, and SH, or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

wherein A<sup>-</sup> is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation;

$R^7$ ,  $R^{7A}$ ,  $R^8$ , and  $R^{8A}$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $R^X$  and  $R^{XA}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heterocycle, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituent selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,

$\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ , and  $\text{C}(\text{O})\text{OM}$ ,

wherein in  $\text{R}^*$  and  $\text{R}^{\text{A}}$ , one or more carbons are optionally replaced by O,  $\text{NR}^{13}$ ,  $\text{N}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^{13}\text{A}^-$ ,  $\text{PR}^{13}$ ,  $\text{P}(\text{O})\text{R}^{13}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{P}(\text{O})\text{R}^9$ ;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S}(\text{O})\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ , CN, OM,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{P}(\text{O})\text{R}^{13}\text{R}^{14}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  $\text{P}(\text{OR}^{13})\text{OR}^{14}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ .

$\text{R}^{19}$  is selected from the group consisting of alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide, polypeptide, wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide polypeptide, can optionally have one or more carbon replaced by O,  $\text{NR}^7$ ,  $\text{N}^+\text{R}^7\text{R}^8$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^7\text{R}^8$ ,  $\text{PR}^7$ ,  $\text{P}^+\text{R}^7\text{R}^8$ , phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl,

wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl,

polyalkoxy diyl, carbohydrate, amino acid, peptide, and polypeptide can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ;

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,  $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  $P^+R^7R^8A^-$ , or phenylene.

122. A compound of claim 121, wherein  $R^1$ ,  $R^{1A}$ ,  $R^2$ , and  $R^{2A}$  are independently selected from the group consisting of H and alkyl.

123. A compound of claim 122, wherein  $R^1$ ,  $R^{1A}$ ,  $R^2$ , and  $R^{2A}$  are independently selected from the group

consisting of H and C<sub>1</sub>-C<sub>10</sub> alkyl.

124. A compound of claim 123, wherein said alkyl is a C<sub>2</sub>-C<sub>4</sub> alkyl.

5

125. A compound of claim 124, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>, and R<sup>2A</sup> are independently C<sub>2</sub>-C<sub>4</sub> alkyl.

10

126. A compound of claim 125, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>, and R<sup>2A</sup> are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.

15

127. A compound of claim 125, wherein R<sup>3</sup>, R<sup>3A</sup>, R<sup>4</sup>, and R<sup>4A</sup> are independently selected from the group consisting of H and OR<sup>9</sup>.

128. A compound of claim 127, wherein R<sup>9</sup> is H.

20

129. A compound of claim 121, wherein R<sup>7</sup>, R<sup>7A</sup>, R<sup>8</sup>, and R<sup>8A</sup> are H.

130. A compound of claim 121, wherein d and e are independently 1 or 2.

25

131. A compound of claim 130, wherein d and e are both 2.

30

132. A compound of claim 121, wherein one or more R<sup>x</sup> and one or more R<sup>xA</sup> are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR<sup>13</sup>, NR<sup>13</sup>R<sup>14</sup>, NR<sup>13</sup>NR<sup>14</sup>R<sup>15</sup>, N<sup>+</sup>R<sup>9</sup>R<sup>11</sup>R<sup>12</sup>A<sup>-</sup>, SR<sup>13</sup>, S<sup>+</sup>R<sup>13</sup>R<sup>14</sup>, CO<sub>2</sub>R<sup>13</sup>, NR<sup>14</sup>C(O)R<sup>13</sup>, and NR<sup>14</sup>C(O)R<sup>13</sup>, wherein alkyl, aryl, cycloalkyl, heterocycle,

polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ .

133. A compound of claim 121, wherein one or more  $R^Y$  and one or more  $R^{YA}$  are independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide,  $NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ , and  $OR^{13}$ ,

wherein alkyl and polyether can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

134. A compound of claim 121, wherein  $R^{19}$  is selected from the group consisting of alkane diyl, polyalkane diyl, alkoxy diyl, and polyalkoxy diyl, wherein alkane diyl and polyalkane diyl can optionally have one or more carbon replaced by O,  $NR^7$ ,  $N^+R^7R^8$ , S, SO,  $SO_2$ ,  $S^+R^7R^8$ ,  $PR^7$ ,  $P^+R^7R^8$ , or phenylene.

SRL 6071  
(C-3021/2)

566

135. A compound of claim 134, wherein  $R^{19}$  is selected from the group consisting of alkoxy diyl and polyalkoxydiyl wherein one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}$ , S, SO, SO<sub>2</sub>,  $S^+R^9R^{10}$ ,  $PR^9$ ,  $P^+R^9R^{10}$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, or polyalkyl.

136. A compound of claim 135, wherein  $R^1$ ,  $R^{1A}$ ,  $R^2$ , and  $R^{2A}$  are independently selected from the group consisting of H and alkyl.

137. A compound of claim 136, wherein  $R^3$ ,  $R^{3A}$ ,  $R^4$ , and  $R^{4A}$  are independently selected from the group consisting of H and  $OR^9$ .

138. A compound of claim 137, wherein  $R^9$  is H.


139. A compound of claim 138, wherein  $R^7$ ,  $R^{7A}$ ,  $R^8$ , and  $R^{8A}$  are each H.

140. A compound of claim 139, wherein d and e are independently 1 or 2.

SRL 6071  
(C-3021/2)

567

141. A compound of claim 140, having the formula:



PEG = 3400 molecular weight polyethylene glycol polymer chain


5

142. A compound of formula (V)



SRL 6071  
(C-3021/2)

568



wherein :

q is an integer from 0 to 4;

r is an integer from 0 to 3;

d and e are independently integers from 0 to 2;

t is an integer from 0 to 4;

u is an integer from 0 to 5;

$R^1$ ,  $R^{1A}$ ,  $R^2$ , and  $R^{2A}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituent selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^{11}A^-$ ,  $SR^9$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, polyalkyl, aryl, and cycloalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,

$N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^w$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which they are attached form C<sub>3</sub>-C<sub>10</sub> cycloalkylidene, or

$R^{1A}$  and  $R^{2A}$  taken together with the carbon to which they are attached form C<sub>3</sub>-C<sub>10</sub> cycloalkylidene;

$R^3$ ,  $R^{3A}$ ,  $R^4$ , and  $R^{4A}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, and SO<sub>3</sub>R<sup>9</sup>, wherein  $R^9$  and  $R^{10}$  are as defined above; or

$R^3$  and  $R^4$  together form =O, =NOR<sup>11</sup>, =S, =NNR<sup>11</sup>R<sup>12</sup>, =NR<sup>9</sup>, or =CR<sup>11</sup>R<sup>12</sup>, or

$R^{3A}$  and  $R^{4A}$  together form =O, =NOR<sup>11</sup>, =S, =NNR<sup>11</sup>R<sup>12</sup>, =NR<sup>9</sup>, or =CR<sup>11</sup>R<sup>12</sup>,

wherein  $R^{11}$  and  $R^{12}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, SO<sub>3</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, CN, halogen, oxo, and CONR<sup>9</sup>R<sup>10</sup>,

wherein  $R^9$  and  $R^{10}$  are as defined above, provided that both  $R^3$  and  $R^4$  cannot be OH, NH<sub>2</sub>, and SH, or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

wherein  $A^-$  is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation;

$R^7$ ,  $R^{7A}$ ,  $R^8$ , and  $R^{8A}$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $R^X$  and  $R^{XA}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heterocycle, polyether, quaternary heterocycle, quaternary

heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,

$CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary

heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituent selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,

SO<sub>2</sub>OM, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, PO(OR<sup>16</sup>)OR<sup>17</sup>, and C(O)OM,

wherein in R<sup>x</sup> and R<sup>xa</sup>, one or more carbons are optionally replaced by O, NR<sup>13</sup>, N<sup>+</sup>R<sup>13</sup>R<sup>14</sup>A<sup>-</sup>, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>13</sup>A<sup>-</sup>, PR<sup>13</sup>, P(O)R<sup>13</sup>, P<sup>+</sup>R<sup>13</sup>R<sup>14</sup>A<sup>-</sup>, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR<sup>9</sup>, N<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>9</sup>A<sup>-</sup>, PR<sup>9</sup>, P<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, or P(O)R<sup>9</sup>;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR<sup>13</sup>, NR<sup>13</sup>R<sup>14</sup>, SR<sup>13</sup>, S(O)R<sup>13</sup>, SO<sub>2</sub>R<sup>13</sup>, SO<sub>3</sub>R<sup>13</sup>, NR<sup>13</sup>OR<sup>14</sup>, NR<sup>13</sup>NR<sup>14</sup>R<sup>15</sup>, NO<sub>2</sub>, CO<sub>2</sub>R<sup>13</sup>, CN, OM, SO<sub>2</sub>OM, SO<sub>2</sub>NR<sup>13</sup>R<sup>14</sup>, C(O)NR<sup>13</sup>R<sup>14</sup>, C(O)OM, COR<sup>13</sup>, P(O)R<sup>13</sup>R<sup>14</sup>, P<sup>+</sup>R<sup>13</sup>R<sup>14</sup>R<sup>15</sup>A<sup>-</sup>, P(OR<sup>13</sup>)OR<sup>14</sup>, S<sup>+</sup>R<sup>13</sup>R<sup>14</sup>A<sup>-</sup>, and N<sup>+</sup>R<sup>9</sup>R<sup>11</sup>R<sup>12</sup>A<sup>-</sup>,

R<sup>19</sup> is selected from the group consisting of alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide, polypeptide, wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide polypeptide, can optionally have one or more carbon replaced by O, NR<sup>7</sup>, N<sup>+</sup>R<sup>7</sup>R<sup>8</sup>, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>7</sup>R<sup>8</sup>, PR<sup>7</sup>, P<sup>+</sup>R<sup>7</sup>R<sup>8</sup>, phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl,

wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl,

SRL 6071  
(C-3021/2)

572

polyalkoxy diyl, carbohydrate, amino acid, peptide, and  
polypeptide can be substituted with one or more  
substituent groups independently selected from the  
group consisting of alkyl, alkenyl, alkynyl, polyalkyl,  
5 polyether, aryl, haloalkyl, cycloalkyl, heterocycle,  
arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  
 $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  
 $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  
 $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  
10  $N^+R^9R^{11}R^{12}A^-$ ;

wherein one or more  $R^y$  and  $R^{yA}$  are independently  
selected from the group consisting of H, alkyl,  
alkenyl, alkynyl, aryl, cycloalkyl, heterocycle,  
quaternary heterocycle,  $OR^9$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  
15  $SO_3R^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl,  
and heterocycle can be substituted with one or more  
substituent groups independently selected from the  
group consisting of alkyl, alkenyl, alkynyl, polyalkyl,  
20 polyether, aryl, haloalkyl, cycloalkyl, heterocycle,  
arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  
 $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  
 $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  
 $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  
25  $N^+R^9R^{11}R^{12}A^-$ ,

wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can be further substituted with one or more substituent  
groups selected from the group consisting of  $OR^7$ ,  
30  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,

CONR<sup>7</sup>R<sup>8</sup>, N<sup>+</sup>R<sup>7</sup>R<sup>8</sup>R<sup>9</sup>A<sup>-</sup>, alkyl, alkenyl, alkynyl, aryl,  
cycloalkyl, heterocycle, arylalkyl, quaternary  
heterocycle, quaternary heteroaryl, P(O)R<sup>7</sup>R<sup>8</sup>, P<sup>+</sup>R<sup>7</sup>R<sup>8</sup>A<sup>-</sup>,  
and P(O)(OR<sup>7</sup>)OR<sup>8</sup>, and

5        wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can optionally have one or more carbons replaced by O,  
NR<sup>7</sup>, N<sup>+</sup>R<sup>7</sup>R<sup>8</sup>A<sup>-</sup>, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>7</sup>A<sup>-</sup>, PR<sup>7</sup>, P(O)R<sup>7</sup>,  
P<sup>+</sup>R<sup>7</sup>R<sup>8</sup>A<sup>-</sup>, or phenylene.

10        143. A compound of claim 142, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently selected from the group  
consisting of H and alkyl.

15        144. A compound of claim 143, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently selected from the group  
consisting of H and C<sub>1</sub>-C<sub>10</sub> alkyl.

20        145. A compound of claim 144, wherein said alkyl  
is a C<sub>2</sub>-C<sub>7</sub> alkyl.

      146. A compound of claim 145, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently C<sub>2</sub>-C<sub>4</sub> alkyl.

25        147. A compound of claim 146, wherein R<sup>1</sup>, R<sup>1A</sup>, R<sup>2</sup>,  
and R<sup>2A</sup> are independently selected from the group  
consisting of ethyl, n-propyl, n-butyl, and isobutyl.

30        148. A compound of claim 142, wherein R<sup>3</sup>, R<sup>3A</sup>, R<sup>4</sup>,  
and R<sup>4A</sup> are independently selected from the group  
consisting of H and OR<sup>9</sup>.

      149. A compound of claim 148, wherein R<sup>9</sup> is H.

150. A compound of claim 142, wherein  $R^7$ ,  $R^{7A}$ ,  $R^8$ , and  $R^{8A}$  are H.

5 151. A compound of claim 142, wherein d and e are independently 1 or 2.

152. A compound of claim 151, wherein d and e are both 2.

10 153. A compound of claim 142, wherein one or more  $R^X$  and one or more  $R^{xA}$  are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^{13}$ ,  $S^+R^{13}R^{14}$ ,  $CO_2R^{13}$ ,  $NR^{14}C(O)R^{13}$ , and  $NR^{14}C(O)R^{13}$ ,

15 wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

20 wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

25 wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ .

30

154. A compound of claim 142, wherein one or more  $R^Y$  and one or more  $R^{YA}$  are independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide,  $NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ , and  $OR^{13}$ ,

wherein alkyl and polyether can be further substituted with  $SO_3R^9$ ,  $N^+R^9R^{11}R^{12}A^-$ , and quaternary heteroaryl.

155. A compound of claim 142, wherein  $R^{19}$  is selected from the group consisting of alkane diyl, polyalkane diyl, alkoxy diyl, and polyalkoxy diyl, wherein alkane diyl and polyalkane diyl can optionally have one or more carbon replaced by O,  $NR^7$ ,  $N^+R^7R^8$ , S, SO,  $SO_2$ ,  $S^+R^7R^8$ ,  $PR^7$ ,  $P^+R^7R^8$ , or phenylene.

156. A compound of claim 155, wherein  $R^{19}$  is selected from the group consisting of alkoxy diyl and polyalkoxydiyl wherein one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}$ , S, SO,  $SO_2$ ,  $S^+R^9R^{10}$ ,  $PR^9$ ,  $P^+R^9R^{10}$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, or polyalkyl.

157. A compound of claim 156, wherein  $R^1$ ,  $R^{1A}$ ,  $R^2$ , and  $R^{2A}$  are independently selected from the group consisting of H and alkyl.

158. A compound of claim 157, wherein  $R^3$ ,  $R^{3A}$ ,  $R^4$ , and  $R^{4A}$  are independently selected from the group consisting of H and  $OR^9$ .

159. A compound of claim 158, wherein  $R^9$  is H.




SRL 6071  
(C-3021/2)

576

160. A compound of claim 159, wherein  $R^7$ ,  $R^{7A}$ ,  $R^8$ , and  $R^{8A}$  are each H.

5 161. A compound of claim 160, wherein d and e are independently 1 or 2.

162. A compound of claim 161, having the formula:



PEG = 3400 molecular weight polyethylene glycol polymer chain

10 163. A pharmaceutical composition comprising an anti-hyperlipidemic condition effective amount of a compound of formula (I) of claim 1, and a pharmaceutically acceptable carrier.

15 164. A pharmaceutical composition comprising an anti-atherosclerotic effective amount of a compound of

a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent

5 groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ , CN, oxo,

$CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary

heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,

15  $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  $P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, aryl, arylalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary heterocycle, quaternary heteroaryl, and

20 quaternary heteroarylalkyl, wherein alkyl, alkenyl, alkynyl, arylalkyl, heterocycle, and polyalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ ,  $P(O)R^9$ , phenylene, carbohydrate, amino acid, peptide, or polypeptide, and

25  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are optionally substituted with one or more groups selected from the group consisting of sulfoalkyl, quaternary heterocycle, quaternary heteroaryl,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,

30

$\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{R}^{11}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , and  $\text{C}(\text{O})\text{OM}$ ,

wherein  $\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected from the substituents constituting  $\text{R}^9$  and  $\text{M}$ ; or

$\text{R}^{14}$  and  $\text{R}^{15}$ , together with the nitrogen atom to which they are attached, form a cyclic ring; and

$\text{R}^6$  is hydroxy; and

$\text{R}^7$  and  $\text{R}^8$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $\text{R}^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary

heteroaryl,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S}(\text{O})\text{R}^{13}$ ,  $\text{S}(\text{O})_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ ,  $\text{CN}$ ,  $\text{OM}$ ,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$ ,  $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{14}\text{C}(\text{O})\text{R}^{13}$ ,  $\text{C}(\text{O})\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{OR}^{18}$ ,  $\text{S}(\text{O})_n\text{NR}^{18}$ ,  $\text{NR}^{13}\text{R}^{18}$ ,  $\text{NR}^{18}\text{OR}^{14}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{P}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary

heteroaryl can be further substituted with  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ ,  $\text{CN}$ , halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ ,  $\text{P}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{C}(\text{O})\text{OM}$ , and

wherein  $\text{R}^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle,

heteroaryl, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

wherein in  $R^X$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ ;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,

provided that both  $R^5$  and  $R^6$  cannot be hydrogen, OH, or SH;

SRL 6071  
(C-3021/2)


583

provided that when  $R^5$  is phenyl, only one of  $R^1$  or  $R^2$  is H; or

a pharmaceutically acceptable salt, solvate, or prodrug thereof.

5

170. A compound of formula I:



wherein:

$q$  is 1 or 2;


$n$  is 2;

$R^1$  and  $R^2$  are each alkyl;

$R^3$  is hydroxy;

$R^4$  and  $R^6$  are hydrogen;

$R^5$  has the formula (II)



(II)

wherein  $t$  is an integer from 0 to 5;

one or more  $R^y$  are  $OR^{13}$ ;

$R^{13}$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, aryl, arylalkyl, cycloalkyl, heterocycle, heteroaryl,

20

SRL 6071  
(C-3021/2)

584

quaternary heterocycle, quaternary heteroaryl, and  
quaternary heteroarylalkyl;

5        said  $R^{13}$  alkyl, alkenyl, alkynyl, arylalkyl,  
heterocycle, and polyalkyl groups optionally have one  
or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S,  $SO$ ,  $SO_2$ ,  
 $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ ,  $P(O)R^9$ , phenylene, carbohydrate,  
amino acid, peptide, or polypeptide;

10         $R^{13}$  is optionally substituted with one or more  
groups selected from the group consisting of  
sulfoalkyl, quaternary heterocycle, quaternary  
heteroaryl,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  
 $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  
 $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S^+R^9R^{10}A^-$ , and  $C(O)OM$ ,

15        wherein  $A^-$  is a pharmaceutically acceptable anion,  
and M is a pharmaceutically acceptable cation,

$R^9$  and  $R^{10}$  are independently selected from the  
group consisting of H, alkyl, alkenyl, alkynyl,  
cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl,  
arylalkyl, and alkylammoniumalkyl;

20         $R^{11}$  and  $R^{12}$  are independently selected from the  
group consisting of H, alkyl, alkenyl, alkynyl, aryl,  
arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle,  
carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl,  
25         $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen,  
oxo, and  $CONR^9R^{10}$ , wherein  $R^9$  and  $R^{10}$  are as defined  
above, provided that both  $R^3$  and  $R^4$  cannot be OH,  $NH_2$ ,  
and SH; or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon  
atom to which they are attached form a cyclic ring; and

30         $R^{16}$  and  $R^{17}$  are independently selected from the  
substituents constituting  $R^9$  and M;

$R^7$  and  $R^8$  are hydrogen; and

SRL 6071  
(C-3021/2)

585

one or more  $R^*$  are independently selected from the group consisting of alkoxy, alkylamino and dialkylamino; or

5 a pharmaceutically acceptable salt, solvate, or prodrug thereof.

171. A compound of claim 170 wherein  $R^1$  and  $R^2$  are each n-butyl.

10 172. A compound of claim 171 wherein t is 1,  $R^y$  is  $OR^{13}$ , and  $R^{13}$  is as defined in claim 170.

15 173. A compound of claim 172 wherein one or more  $R^*$  are independently selected from methoxy and dimethylamino.

174. A compound of claim 172 wherein  $R^*$  is dimethylamino.

20 175. A compound of claim 172 wherein:  
t is 1;  
 $R^y$  is para- $OR^{13}$ ; and  
 $R^{13}$  is as defined in claim 170.


25 176. A compound of claim 172 wherein:  
t is 1;  
 $R^y$  is meta- $OR^{13}$ ; and  
 $R^{13}$  is as defined in claim 170.

30 177. A compound of claim 172 having the 4R,5R configuration.


178. A compound of claim 170 having the structural formula:

SRL 6071  
(C-3021/2)

586




179. A compound of claim 170 having the  
5 structural formula:



180. A compound of formula (I):





wherein:

q is an integer from 1 to 4;

n is an integer from 0 to 2;

$R^1$  and  $R^2$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^wA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^w$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,

ammoniumalkyl, arylalkyl, carboxyalkyl,  
carboxyheteroaryl, carboxyheterocycle,  
carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,  
heterocyclalkyl, and alkylammoniumalkyl; or

5  $R^1$  and  $R^2$  taken together with the carbon to which  
they are attached form  $C_3$ - $C_{10}$  cycloalkyl;

$R^3$  and  $R^4$  are independently selected from the  
group consisting of H, alkyl, alkenyl, alkynyl,  
acyloxy, aryl, heterocycle,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  
10  $SO_2R^9$ , and  $SO_3R^9$ , wherein  $R^9$  and  $R^{10}$  are as defined  
above; or

$R^3$  and  $R^4$  together form  $=O$ ,  $=NOR^{11}$ ,  $=S$ ,  $=NNR^{11}R^{12}$ ,  
 $=NR^9$ , or  $=CR^{11}R^{12}$ ,

wherein  $R^{11}$  and  $R^{12}$  are independently selected  
15 from the group consisting of H, alkyl, alkenyl,  
alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl,  
heterocycle, carboxyalkyl, carboalkoxyalkyl,  
cycloalkyl, cyanoalkyl,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  
 $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,  
20 wherein  $R^9$  and  $R^{10}$  are as defined above, provided that  
both  $R^3$  and  $R^4$  cannot be OH,  $NH_2$ , and SH, or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon  
atom to which they are attached form a cyclic ring;

$R^5$  is aryl substituted with one or more  $OR^{13a}$ ,

25 wherein  $R^{13a}$  is selected from the group consisting  
of polyether, aryl, alkylarylalkyl,  
alkylheteroarylalkyl, alkylheterocyclalkyl,  
heterocyclalkyl, heteroarylalkyl, quaternary  
heterocyclalkyl, alkylammoniumalkyl, and  
30 carboxyalkylaminocarbonylalkyl,

$R^{13a}$  is optionally substituted with one or more groups selected from the group consisting of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl, heterocycle, heteroaryl, sulfoalkyl, quaternary heterocycle, quaternary heteroaryl, quaternary heterocyclylalkyl, quaternary heteroarylalkyl, guanidinyl,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S^+R^9R^{10}A^-$ , and  $C(O)OM$ ,

wherein  $A^-$  is an pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation,

wherein  $R^{16}$  and  $R^{17}$  are independently selected from the substituents constituting  $R^9$  and M; and

$R^6$  is selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle,  $OR^{30}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $NR^{13}C(O)R^{14}$ ,  $NR^{13}C(O)NR^{14}R^{15}$ ,  $NR^{13}CO_2R^{14}$ ,  $OC(O)R^{13}$ ,  $OC(O)NR^{13}R^{14}$ ,  $NR^{13}SOR^{14}$ ,  $NR^{13}SO_2R^{14}$ ,  $NR^{13}SONR^{14}R^{15}$ ,

$\text{NR}^{13}\text{SO}_2\text{NR}^{14}\text{R}^{15}$ ,  $\text{P}(\text{O})\text{R}^{13}\text{R}^{14}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  $\text{P}(\text{OR}^{13})\text{OR}^{14}$ ,  
 $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,

wherein:

$\text{A}^-$  is a pharmaceutically acceptable anion and M is  
 5 a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl,  
 polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
 can be further substituted with one or more substituent  
 groups selected from the group consisting of  $\text{OR}^7$ ,  
 10  $\text{NR}^7\text{R}^8$ ,  $\text{SR}^7$ ,  $\text{S}(\text{O})\text{R}^7$ ,  $\text{SO}_2\text{R}^7$ ,  $\text{SO}_3\text{R}^7$ ,  $\text{CO}_2\text{R}^7$ , CN, oxo,  
 $\text{CONR}^7\text{R}^8$ ,  $\text{N}^+\text{R}^7\text{R}^8\text{R}^9\text{A}^-$ , alkyl, alkenyl, alkynyl, aryl,  
 cycloalkyl, heterocycle, arylalkyl, quaternary  
 heterocycle, quaternary heteroaryl,  $\text{P}(\text{O})\text{R}^7\text{R}^8$ ,  
 $\text{P}^+\text{R}^7\text{R}^8\text{R}^9\text{A}^-$ , and  $\text{P}(\text{O})(\text{OR}^7)\text{OR}^8$ , and

15 wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
 polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
 can optionally have one or more carbons replaced by O,  
 $\text{NR}^7$ ,  $\text{N}^+\text{R}^7\text{R}^8\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^7\text{A}^-$ ,  $\text{PR}^7$ ,  $\text{P}(\text{O})\text{R}^7$ ,  
 $\text{P}^+\text{R}^7\text{R}^8\text{A}^-$ , or phenylene, and  $\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are  
 20 independently selected from the group consisting of  
 hydrogen, alkyl, alkenyl, alkynyl, polyalkyl,  
 polyether, aryl, arylalkyl, alkylarylalkyl,  
 alkylheteroarylalkyl, alkylheterocyclalkyl,  
 cycloalkyl, heterocycle, heteroaryl, quaternary  
 25 heterocycle, quaternary heteroaryl, heterocyclalkyl,  
 heteroarylalkyl, quaternary heterocyclalkyl,  
 quaternary heteroarylalkyl, alkylammoniumalkyl, and  
 carboxyalkylaminocarbonylalkyl,

30 wherein alkyl, alkenyl, alkynyl, arylalkyl,  
 heterocycle, and polyalkyl optionally have one or more  
 carbons replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,

$S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ ,  $P(O)R^9$ , phenylene, carbohydrate, amino acid, peptide, or polypeptide, and

$R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are optionally substituted with one or more groups selected from the group consisting of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl, heterocycle, heteroaryl, sulfoalkyl, quaternary heterocycle, quaternary heteroaryl, quaternary heterocyclalkyl, quaternary heteroaryalkyl, guanidiny,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S^+R^9R^{10}A^-$ , and  $C(O)OM$ ,

wherein  $R^{16}$  and  $R^{17}$  are independently selected from the substituents constituting  $R^9$  and M; or

$R^{13}$  and  $R^{14}$ , together with the nitrogen atom to which they are attached form a mono- or polycyclic heterocycle that is optionally substituted with one or more radicals selected from the group consisting of oxo, carboxy and quaternary salts; or

$R^{14}$  and  $R^{15}$ , together with the nitrogen atom to which they are attached, form a cyclic ring; and

$R^{30}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroaryalkyl, heterocyclalkyl, and alkylammoniumalkyl; and

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $R^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl,

polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

wherein in  $R^X$ , one or more carbons are optionally replaced by  $O$ ,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ ,  $S$ ,  $SO$ ,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid,

peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more  
5 carbons are optionally replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{P}(\text{O})\text{R}^9$ ;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more  
10 groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S}(\text{O})\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ , CN, OM,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{P}(\text{O})\text{R}^{13}\text{R}^{14}$ ,  
15  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  $\text{P}(\text{OR}^{13})\text{OR}^{14}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ , or

a pharmaceutically acceptable salt, solvate, or prodrug thereof.

181. A compound of claim 180 wherein:

$\text{R}^5$  is phenyl substituted with  $\text{OR}^{13a}$ ;

$\text{R}^{13a}$  is independently selected from the group  
25 consisting of polyether, alkylarylalkyl, alkylheteroarylalkyl, alkylheterocyclalkyl, and carboxyalkylaminocarbonylalkyl; and

$\text{R}^{13a}$  is optionally substituted with one or more  
30 groups selected from the group consisting of carboxy, quaternary heterocycle, quaternary heteroaryl, and  $\text{NR}^9\text{R}^{10}$ .

182. A compound of claim 180 wherein n is 1 or 2.

183. A compound of claim 180 wherein  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl.

5

184. A compound of claim 180 wherein  $R^7$  and  $R^8$  are hydrogen.

10

185. A compound of claim 180 wherein  $R^8$  and  $R^4$  are independently selected from the group consisting of hydrogen and  $OR^9$ .

15

186. A compound of claim 180 wherein  $R^3$  is hydrogen and  $R^4$  is hydroxy.

20

187. A compound of claim 180 wherein one or more  $R^x$  are independently selected from the group consisting of  $OR^{13}$  and  $NR^{13}R^{14}$ .

188. A compound of claim 180 wherein one or more  $R^x$  are independently selected from methoxy and dimethylamino.

25

189. A compound of claim 180 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen and alkyl.

30

190. A compound of claim 180 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting alkyl.

191. A compound of claim 180 wherein  $R^1$  and  $R^2$  are the same alkyl.

35

192. A compound of claim 180 wherein  $R^1$  and  $R^2$  are each n-butyl.




SRL 6071  
(C-3021/2)


595

193. A compound of claim 180 wherein  
n is 1 or 2;  
R<sup>1</sup> and R<sup>2</sup> are n-butyl;  
R<sup>3</sup> and R<sup>6</sup> are hydrogen;  
R<sup>4</sup> is hydroxy;  
R<sup>7</sup> and R<sup>8</sup> are hydrogen; and  
one or more R\* are independently selected from  
methoxy and dimethylamino.

194. A compound of claim 180 having the structural  
formula:




195. A compound of claim 180 having the structural  
formula:




196. A compound of claim 180 having the structural  
formula:

SRL 6071  
(C-3021/2)

596




197. A compound of claim 180 having the structural formula:



5


198. A compound of claim 180 having the structural formula:




199. A compound of claim 180 having the structural formula:

SRL 6071  
(C-3021/2)


597



200. A compound of claim 180 having the structural formula:



201. A compound of formula (I):



wherein:

q is an integer from 1 to 4;

n is an integer from 0 to 2;

R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and

cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^WA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^W$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclylalkyl, and alkylammoniumalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which they are attached form  $C_3$ - $C_{10}$  cycloalkyl;

$R^3$  and  $R^4$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ , wherein  $R^9$  and  $R^{10}$  are as defined above; or

$R^3$  and  $R^4$  together form  $=O$ ,  $=NOR^{11}$ ,  $=S$ ,  $=NNR^{11}R^{12}$ ,  $=NR^9$ , or  $=CR^{11}R^{12}$ ,

wherein  $R^{11}$  and  $R^{12}$  are independently selected

from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, SO<sub>3</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, CN, halogen, oxo, and CONR<sup>9</sup>R<sup>10</sup>,  
5 wherein R<sup>9</sup> and R<sup>10</sup> are as defined above, provided that both R<sup>3</sup> and R<sup>4</sup> cannot be OH, NH<sub>2</sub>, and SH, or

R<sup>11</sup> and R<sup>12</sup> together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

10 R<sup>5</sup> is aryl substituted with one or more OR<sup>13b</sup>,

wherein R<sup>13b</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, arylalkyl, alkylarylalkyl, alkylheteroarylalkyl, alkylheterocyclalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary heterocycle, quaternary heteroaryl, heterocyclalkyl, heteroarylalkyl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, alkylammoniumalkyl, and carboxyalkylaminocarbonylalkyl,

20 R<sup>13b</sup> is substituted with one or more groups selected from the group consisting of hydroxy, amino, sulfo, carboxy, carboxyalkyl, heterocycle, heteroaryl, sulfoalkyl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, or guanidiny, and

25 R<sup>6</sup> is selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, OR<sup>30</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, and SO<sub>3</sub>R<sup>9</sup>,

30 wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more

substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $NR^{13}C(O)R^{14}$ ,  $NR^{13}C(O)NR^{14}R^{15}$ ,  $NR^{13}CO_2R^{14}$ ,  $OC(O)R^{13}$ ,  $OC(O)NR^{13}R^{14}$ ,  $NR^{13}SOR^{14}$ ,  $NR^{13}SO_2R^{14}$ ,  $NR^{13}SONR^{14}R^{15}$ ,  $NR^{13}SO_2NR^{14}R^{15}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,

wherein:

$A^-$  is a pharmaceutically acceptable anion and  $M$  is a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by  $O$ ,  $NR^7$ ,  $N^+R^7R^8A^-$ ,  $S$ ,  $SO$ ,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  $P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl,

polyether, aryl, arylalkyl, alkylarylalkyl,  
alkylheteroarylalkyl, alkylheterocyclalkyl,  
cycloalkyl, heterocycle, heteroaryl, quaternary  
heterocycle, quaternary heteroaryl, heterocyclalkyl,  
5 heteroarylalkyl, quaternary heterocyclalkyl,  
quaternary heteroarylalkyl, alkylammoniumalkyl, and  
carboxyalkylaminocarbonylalkyl,

wherein alkyl, alkenyl, alkynyl, arylalkyl,  
heterocycle, and polyalkyl optionally have one or more  
10 carbons replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  
 $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ ,  $\text{P}(\text{O})\text{R}^9$ , phenylene, carbohydrate,  
amino acid, peptide, or polypeptide, and

$\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are optionally substituted with  
one or more groups selected from the group consisting  
15 of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl,  
heterocycle, heteroaryl, sulfoalkyl, quaternary  
heterocycle, quaternary heteroaryl, quaternary  
heterocyclalkyl, quaternary heteroarylalkyl,  
guanidinyl,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
20  $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ , CN, halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_2\text{OM}$ ,  
 $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{R}^{11}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , and  
 $\text{C}(\text{O})\text{OM}$ ,

wherein  $\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected  
from the substituents constituting  $\text{R}^9$  and M; or

25  $\text{R}^{13}$  and  $\text{R}^{14}$ , together with the nitrogen atom to  
which they are attached form a mono- or polycyclic  
heterocycle that is optionally substituted with one or  
more radicals selected from the group consisting of  
oxo, carboxy and quaternary salts; or

30  $\text{R}^{14}$  and  $\text{R}^{15}$ , together with the nitrogen atom to  
which they are attached, form a cyclic ring; and

$R^{30}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclalkyl, and alkylammoniumalkyl; and

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $R^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl,



heterocycle, heteroaryl, alkyl, quaternary heterocycle,  
and quaternary heteroaryl optionally are substituted  
with one or more substituents selected from the group  
consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  
5  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  
 $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

wherein in  $R^x$ , one or more carbons are optionally  
replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  
 $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid,  
10 peptide, polypeptide, carbohydrate, polyether, or  
polyalkyl,

wherein in said polyalkyl, phenylene, amino acid,  
peptide, polypeptide, and carbohydrate, one or more  
carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ ,  
15 S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ ;

wherein quaternary heterocycle and quaternary  
heteroaryl are optionally substituted with one or more  
groups selected from the group consisting of alkyl,  
alkenyl, alkynyl, polyalkyl, polyether, aryl,  
20 haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,  
oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  
 $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  
 $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  
 $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  
25  $N^+R^9R^{11}R^{12}A^-$ , or

a pharmaceutically acceptable salt, solvate, or  
prodrug thereof.

202. A compound of claim 201 wherein:  
30  $R^5$  is phenyl substituted with  $OR^{13b}$ ;

$R^{13b}$  is independently selected from the group consisting of alkyl, quaternary heteroarylalkyl, and quaternary heterocyclylalkyl; and

5  $R^{13b}$  is substituted with one or more groups selected from the group consisting of hydroxy, heterocycle, heteroaryl, and guanidiny1.

203. A compound of claim 201 wherein  $n$  is 1 or 2.

10 204. A compound of claim 201 wherein  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl.

15 205. A compound of claim 201 wherein  $R^7$  and  $R^8$  are hydrogen.

20 206. A compound of claim 201 wherein  $R^3$  and  $R^4$  are independently selected from the group consisting of hydrogen and  $OR^9$ .

207. A compound of claim 201 wherein  $R^3$  is hydrogen and  $R^4$  is hydroxy.

25 208. A compound of claim 201 wherein one or more  $R^x$  are independently selected from the group consisting of  $OR^{13}$  and  $NR^{13}R^{14}$ .

30 209. A compound of claim 201 wherein one or more  $R^x$  are independently selected from methoxy and dimethylamino.

35 210. A compound of claim 201 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen and alkyl.


211. A compound of claim 201 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting alkyl.

212. A compound of claim 201 wherein  $R^1$  and  $R^2$  are the same alkyl.

213. A compound of claim 201 wherein  $R^1$  and  $R^2$  are each n-butyl.

214. A compound of claim 201 wherein  
n is 1 or 2;  
 $R^1$  and  $R^2$  are n-butyl;  
 $R^3$  and  $R^6$  are hydrogen;  
 $R^4$  is hydroxy;  
 $R^7$  and  $R^8$  are hydrogen; and  
one or more  $R^x$  are independently selected from methoxy and dimethylamino.


215. A compound of claim 201 having the structural formula:




216. A compound of claim 201 having the structural formula:

SRL 6071  
(C-3021/2)


606



217. A compound of claim 201 having the structural formula:



218. A compound of formula (I):



(I)

wherein:

$q$  is an integer from 1 to 4;

$n$  is an integer from 0 to 2;

$R^1$  and  $R^2$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more  
 5 substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^WA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl  
 10 optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^W$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclylalkyl, and alkylammoniumalkyl; or  
 15

$R^1$  and  $R^2$  taken together with the carbon to which they are attached form  $C_3$ - $C_{10}$  cycloalkyl;  
 20

$R^3$  and  $R^4$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ , wherein  $R^9$  and  $R^{10}$  are as defined  
 25 above; or

$R^3$  and  $R^4$  together form  $=O$ ,  $=NOR^{11}$ ,  $=S$ ,  $=NNR^{11}R^{12}$ ,  $=NR^9$ , or  $=CR^{11}R^{12}$ ,

wherein  $R^{11}$  and  $R^{12}$  are independently selected from the group consisting of H, alkyl, alkenyl,  
 30

alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl,  
heterocycle, carboxyalkyl, carboalkoxyalkyl,  
cycloalkyl, cyanoalkyl,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,  
 $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,  
5 wherein  $R^9$  and  $R^{10}$  are as defined above, provided that  
both  $R^3$  and  $R^4$  cannot be OH,  $NH_2$ , and SH, or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon  
atom to which they are attached form a cyclic ring;

$R^5$  is aryl substituted with one or more  $OR^{13b}$ ,

10 wherein  $R^{13b}$  is selected from the group consisting  
of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl,  
arylalkyl, alkylarylalkyl, alkylheteroarylalkyl,  
alkylheterocyclalkyl, cycloalkyl, heterocycle,  
heteroaryl, quaternary heterocycle, quaternary  
15 heteroaryl, heterocyclalkyl, heteroarylalkyl,  
quaternary heterocyclalkyl, quaternary  
heteroarylalkyl, alkylammoniumalkyl, and  
carboxyalkylaminocarbonylalkyl,

$R^{13b}$  is substituted with one or more groups  
20 selected from the group consisting of  $OR^{9a}$ ,  $NR^{9a}R^{10}$ ,  
 $N^+R^{9a}R^{11}R^{12}A^-$ ,  $SR^{9a}$ ,  $S(O)R^{9a}$ ,  $SO_2R^{9a}$ ,  $SO_3R^{9a}$ ,  $CO_2R^{9a}$ ,  
 $CONR^{9a}R^{10}$ ,  $SO_2NR^{9a}R^{10}$ ,  $P^+R^{9a}R^{10}R^{11}A^-$ , and  $S^+R^{9a}R^{10}A^-$ ,

wherein  $A^-$  is an pharmaceutically acceptable anion  
and M is a pharmaceutically acceptable cation, and

25 wherein  $R^{9a}$  is selected from the group consisting  
of carboxyalkyl, carboxyheteroaryl, carboxyheterocycle,  
carboalkoxyalkyl, and carboxyalkylamino;

$R^6$  is selected from the group consisting of H,  
alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle,  
30 quaternary heterocycle,  $OR^{30}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and

$\text{SO}_3\text{R}^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, halogen, oxo,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S(O)}\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ ,  $\text{CN}$ ,  $\text{OM}$ ,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{C(O)}\text{NR}^{13}\text{R}^{14}$ ,  $\text{C(O)}\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{NR}^{13}\text{C(O)}\text{R}^{14}$ ,  $\text{NR}^{13}\text{C(O)}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NR}^{13}\text{CO}_2\text{R}^{14}$ ,  $\text{OC(O)}\text{R}^{13}$ ,  $\text{OC(O)}\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{13}\text{SOR}^{14}$ ,  $\text{NR}^{13}\text{SO}_2\text{R}^{14}$ ,  $\text{NR}^{13}\text{SONR}^{14}\text{R}^{15}$ ,  $\text{NR}^{13}\text{SO}_2\text{NR}^{14}\text{R}^{15}$ ,  $\text{P(O)}\text{R}^{13}\text{R}^{14}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  $\text{P(OR}^{13})\text{OR}^{14}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,

wherein:

$\text{A}^-$  is a pharmaceutically acceptable anion and  $\text{M}$  is a pharmaceutically acceptable cation, said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $\text{OR}^7$ ,  $\text{NR}^7\text{R}^8$ ,  $\text{SR}^7$ ,  $\text{S(O)}\text{R}^7$ ,  $\text{SO}_2\text{R}^7$ ,  $\text{SO}_3\text{R}^7$ ,  $\text{CO}_2\text{R}^7$ ,  $\text{CN}$ , oxo,  $\text{CONR}^7\text{R}^8$ ,  $\text{N}^+\text{R}^7\text{R}^8\text{R}^9\text{A}^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $\text{P(O)}\text{R}^7\text{R}^8$ ,  $\text{P}^+\text{R}^7\text{R}^8\text{R}^9\text{A}^-$ , and  $\text{P(O)}(\text{OR}^7)\text{OR}^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,

$NR^7$ ,  $N^+R^7R^8A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  
 $P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are  
independently selected from the group consisting of  
hydrogen, alkyl, alkenyl, alkynyl, polyalkyl,  
5 polyether, aryl, arylalkyl, alkylarylalkyl,  
alkylheteroarylalkyl, alkylheterocyclalkyl,  
cycloalkyl, heterocycle, heteroaryl, quaternary  
heterocycle, quaternary heteroaryl, heterocyclalkyl,  
heteroarylalkyl, quaternary heterocyclalkyl,  
10 quaternary heteroarylalkyl, alkylammoniumalkyl, and  
carboxyalkylaminocarbonylalkyl,

wherein alkyl, alkenyl, alkynyl, arylalkyl,  
heterocycle, and polyalkyl optionally have one or more  
carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  
15  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ ,  $P(O)R^9$ , phenylene, carbohydrate,  
amino acid, peptide, or polypeptide, and

$R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are optionally substituted with  
one or more groups selected from the group consisting  
of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl,  
20 heterocycle, heteroaryl, sulfoalkyl, quaternary  
heterocycle, quaternary heteroaryl, quaternary  
heterocyclalkyl, quaternary heteroarylalkyl,  
guanidinyll,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  
 $SO_2R^9$ ,  $SO_3R^9$ ,  $oxo$ ,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  
25  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S^+R^9R^{10}A^-$ , and  
 $C(O)OM$ ,

wherein  $R^{16}$  and  $R^{17}$  are independently selected  
from the substituents constituting  $R^9$  and M; or  
 $R^{13}$  and  $R^{14}$ , together with the nitrogen atom to  
30 which they are attached form a mono- or polycyclic  
heterocycle that is optionally substituted with one or



SRL 6071  
(C-3021/2)

611

more radicals selected from the group consisting of oxo, carboxy and quaternary salts; or

$R^{14}$  and  $R^{15}$ , together with the nitrogen atom to which they are attached, form a cyclic ring; and

5  $R^{30}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclylalkyl, and alkylammoniumalkyl; and

10  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $R^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  
20  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

25 wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  
30  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

SRL 6071  
(C-3021/2)

612

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl,

5 wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  
10  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

wherein in  $R^x$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or  
15 polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ ;

20 wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,  
25 oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ , or

a pharmaceutically acceptable salt, solvate, or prodrug thereof.

219. A compound of claim 218 wherein:

$R^5$  is phenyl substituted with  $OR^{13b}$ ;

$R^{13b}$  is alkyl; and

$R^{13b}$  is substituted with one or more groups selected from the group consisting of  $OR^{9a}$  and  $NR^{9a}R^{10}$ ; and

$R^{9a}$  is selected from the group consisting of carboxyalkyl, carboxyheteroaryl, and carboxyheterocycle; and

$R^{10}$  is carboxyalkyl.

220. A compound of claim 218 wherein n is 1 or 2.

221. A compound of claim 218 wherein  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl.

222. A compound of claim 218 wherein  $R^7$  and  $R^8$  are hydrogen.

223. A compound of claim 218 wherein  $R^3$  and  $R^4$  are independently selected from the group consisting of hydrogen and  $OR^9$ .

224. A compound of claim 218 wherein  $R^3$  is hydrogen and  $R^4$  is hydroxy.

224. A compound of claim 218 wherein one or more  $R^x$  are independently selected from the group consisting of  $OR^{13}$  and  $NR^{13}R^{14}$ .

226. A compound of claim 218 wherein one or more  $R^x$  are independently selected from methoxy and

SRL 6071  
(C-3021/2)

614

dimethylamino.

5 227. A compound of claim 218 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen and alkyl.

228. A compound of claim 218 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting alkyl.


10 229. A compound of claim 218 wherein  $R^1$  and  $R^2$  are the same alkyl.

230. A compound of claim 218 wherein  $R^1$  and  $R^2$  are each n-butyl.

15 231. A compound of claim 218 wherein  
n is 1 or 2;  
 $R^1$  and  $R^2$  are n-butyl;  
 $R^3$  and  $R^6$  are hydrogen;  
20  $R^4$  is hydroxy;

$R^7$  and  $R^8$  are hydrogen; and  
one or more  $R^i$  are independently selected from methoxy and dimethylamino.

25 232. A compound of claim 218 having the structural formula:




233. A compound of claim 218 having the structural


SRL 6071  
(C-3021/2)

615

formula:



234. A compound of formula (I):



(I)

wherein:

q is an integer from 1 to 4;

n is an integer from 0 to 2;

R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituents selected from the group consisting of OR<sup>9</sup>,

$\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{R}^{\text{w}}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{R}^{11}\text{A}^-$ ,  
 $\text{S}(\text{O})\text{R}^9$ ,  $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ ,  $\text{CO}_2\text{R}^9$ , CN, halogen, oxo, and  
 $\text{CONR}^9\text{R}^{10}$ ,

5        wherein alkyl, alkenyl, alkynyl, alkylaryl,  
 alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl  
 optionally have one or more carbons replaced by O,  $\text{NR}^9$ ,  
 $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or phenylene,

10        wherein  $\text{R}^9$ ,  $\text{R}^{10}$ , and  $\text{R}^{\text{w}}$  are independently selected  
 from the group consisting of H, alkyl, alkenyl,  
 alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
 ammoniumalkyl, arylalkyl, carboxyalkyl,  
 carboxyheteroaryl, carboxyheterocycle,  
 carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,  
 heterocyclylalkyl, and alkylammoniumalkyl; or

15         $\text{R}^1$  and  $\text{R}^2$  taken together with the carbon to which  
 they are attached form  $\text{C}_3\text{-C}_{10}$  cycloalkyl;

20         $\text{R}^3$  and  $\text{R}^4$  are independently selected from the  
 group consisting of H, alkyl, alkenyl, alkynyl,  
 acyloxy, aryl, heterocycle,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
 $\text{SO}_2\text{R}^9$ , and  $\text{SO}_3\text{R}^9$ , wherein  $\text{R}^9$  and  $\text{R}^{10}$  are as defined  
 above; or

$\text{R}^3$  and  $\text{R}^4$  together form  $=\text{O}$ ,  $=\text{NOR}^{11}$ ,  $=\text{S}$ ,  $=\text{NNR}^{11}\text{R}^{12}$ ,  
 $=\text{NR}^9$ , or  $=\text{CR}^{11}\text{R}^{12}$ ,

25        wherein  $\text{R}^{11}$  and  $\text{R}^{12}$  are independently selected  
 from the group consisting of H, alkyl, alkenyl,  
 alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl,  
 heterocycle, carboxyalkyl, carboalkoxyalkyl,  
 cycloalkyl, cyanoalkyl,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
 $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ ,  $\text{CO}_2\text{R}^9$ , CN, halogen, oxo, and  $\text{CONR}^9\text{R}^{10}$ ,

wherein  $R^9$  and  $R^{10}$  are as defined above, provided that both  $R^3$  and  $R^4$  cannot be OH,  $NH_2$ , and SH, or

$R^{11}$  and  $R^{12}$  together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

5  $R^5$  is aryl substituted with one or more  $OR^{13b}$ ,

wherein  $R^{13b}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, arylalkyl, alkylarylalkyl, alkylheteroarylalkyl, alkylheterocyclalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary heterocycle, quaternary heteroaryl, heterocyclalkyl, heteroarylalkyl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, alkylammoniumalkyl, and carboxyalkylaminocarbonylalkyl,

15  $R^{13b}$  is substituted with one or more groups selected from the group consisting of  $NR^9R^{10a}$ ,  $CONR^9R^{10a}$ ,  $SO_2NR^9R^{10a}$ ,  $P^+R^9R^{10a}R^{11}A^-$ , and  $S^+R^9R^{10a}A^-$ ,

wherein  $A^-$  is an pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation,

20 wherein  $R^{10a}$  is selected from the group consisting of carboxyalkyl, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, and heterocyclalkyl; or

$R^6$  is selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle,  $OR^{30}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ ,

25 wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more  
30 substituent groups independently selected from the

SRL 6071  
(C-3021/2)

618

group consisting of alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, heterocycle,  
arylalkyl, quaternary heterocycle, quaternary  
heteroaryl, halogen, oxo,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S(O)}\text{R}^{13}$ ,  
5  $\text{SO}_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ ,  $\text{CN}$ ,  
 $\text{OM}$ ,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  $\text{C(O)}\text{NR}^{13}\text{R}^{14}$ ,  $\text{C(O)}\text{OM}$ ,  $\text{COR}^{13}$ ,  
 $\text{NR}^{13}\text{C(O)}\text{R}^{14}$ ,  $\text{NR}^{13}\text{C(O)}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NR}^{13}\text{CO}_2\text{R}^{14}$ ,  $\text{OC(O)}\text{R}^{13}$ ,  
 $\text{OC(O)}\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{13}\text{SOR}^{14}$ ,  $\text{NR}^{13}\text{SO}_2\text{R}^{14}$ ,  $\text{NR}^{13}\text{SONR}^{14}\text{R}^{15}$ ,  
 $\text{NR}^{13}\text{SO}_2\text{NR}^{14}\text{R}^{15}$ ,  $\text{P(O)}\text{R}^{13}\text{R}^{14}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  $\text{P(OR}^{13})\text{OR}^{14}$ ,  
10  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,

wherein:

$\text{A}^-$  is a pharmaceutically acceptable anion and M is  
a pharmaceutically acceptable cation,  
said alkyl, alkenyl, alkynyl, polyalkyl,  
15 polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can be further substituted with one or more substituent  
groups selected from the group consisting of  $\text{OR}^7$ ,  
 $\text{NR}^7\text{R}^8$ ,  $\text{SR}^7$ ,  $\text{S(O)}\text{R}^7$ ,  $\text{SO}_2\text{R}^7$ ,  $\text{SO}_3\text{R}^7$ ,  $\text{CO}_2\text{R}^7$ ,  $\text{CN}$ , oxo,  
 $\text{CONR}^7\text{R}^8$ ,  $\text{N}^+\text{R}^7\text{R}^8\text{R}^9\text{A}^-$ , alkyl, alkenyl, alkynyl, aryl,  
20 cycloalkyl, heterocycle, arylalkyl, quaternary  
heterocycle, quaternary heteroaryl,  $\text{P(O)}\text{R}^7\text{R}^8$ ,  
 $\text{P}^+\text{R}^7\text{R}^8\text{R}^9\text{A}^-$ , and  $\text{P(O)}(\text{OR}^7)\text{OR}^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
25 can optionally have one or more carbons replaced by O,  
 $\text{NR}^7$ ,  $\text{N}^+\text{R}^7\text{R}^8\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^7\text{A}^-$ ,  $\text{PR}^7$ ,  $\text{P(O)}\text{R}^7$ ,  
 $\text{P}^+\text{R}^7\text{R}^8\text{A}^-$ , or phenylene, and  $\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are  
independently selected from the group consisting of  
hydrogen, alkyl, alkenyl, alkynyl, polyalkyl,  
30 polyether, aryl, arylalkyl, alkylarylalkyl,



alkylheteroarylalkyl, alkylheterocyclalkyl,  
cycloalkyl, heterocycle, heteroaryl, quaternary  
heterocycle, quaternary heteroaryl, heterocyclalkyl,  
heteroarylalkyl, quaternary heterocyclalkyl,  
5 quaternary heteroarylalkyl, alkylammoniumalkyl, and  
carboxyalkylaminocarbonylalkyl,

wherein alkyl, alkenyl, alkynyl, arylalkyl,  
heterocycle, and polyalkyl optionally have one or more  
carbons replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  
10  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ ,  $\text{P}(\text{O})\text{R}^9$ , phenylene, carbohydrate,  
amino acid, peptide, or polypeptide, and

$\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are optionally substituted with  
one or more groups selected from the group consisting  
of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl,  
15 heterocycle, heteroaryl, sulfoalkyl, quaternary  
heterocycle, quaternary heteroaryl, quaternary  
heterocyclalkyl, quaternary heteroarylalkyl,  
guanidiny,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
 $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ , CN, halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_2\text{OM}$ ,  
20  $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{R}^{11}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , and  
 $\text{C}(\text{O})\text{OM}$ ,

wherein  $\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected  
from the substituents constituting  $\text{R}^9$  and M; or  
 $\text{R}^{13}$  and  $\text{R}^{14}$ , together with the nitrogen atom to  
25 which they are attached form a mono- or polycyclic  
heterocycle that is optionally substituted with one or  
more radicals selected from the group consisting of  
oxo, carboxy and quaternary salts; or

$\text{R}^{14}$  and  $\text{R}^{15}$ , together with the nitrogen atom to  
30 which they are attached, form a cyclic ring; and

$\text{R}^{30}$  is selected from the group consisting of alkyl,  
alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,

SRL 6071  
(C-3021/2)

620

ammoniumalkyl, alkylammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclylalkyl, and alkylammoniumalkyl; and

5  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $R^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

25 wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle, and quaternary heteroaryl optionally are substituted

30

with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

5 wherein in  $R^x$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

10 wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ ;

15 wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ , or

25 a pharmaceutically acceptable salt, solvate, or prodrug thereof.

235. A compound of claim 234 wherein:

$R^5$  is phenyl substituted with  $OR^{13b}$ ;

$R^{13b}$  is alkyl; and

30  $R^{13b}$  is substituted with  $NR^9R^{10a}$ ; and

$R^9$  is hydrogen; and  
 $R^{10}$  is heteroarylalkyl.

236. A compound of claim 234 wherein  $n$  is 1 or 2.

5

237. A compound of claim 234 wherein  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl.

10

238. A compound of claim 234 wherein  $R^7$  and  $R^8$  are hydrogen.

15

239. A compound of claim 234 wherein  $R^3$  and  $R^4$  are independently selected from the group consisting of hydrogen and  $OR^9$ .

20

240. A compound of claim 234 wherein  $R^3$  is hydrogen and  $R^4$  is hydroxy.

241. A compound of claim 234 wherein one or more  $R^x$  are independently selected from the group consisting of  $OR^{13}$  and  $NR^{13}R^{14}$ .

25

242. A compound of claim 234 wherein one or more  $R^x$  are independently selected from methoxy and dimethylamino.

30

243. A compound of claim 234 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen and alkyl.

244. A compound of claim 234 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting alkyl.

35

245. A compound of claim 234 wherein  $R^1$  and  $R^2$  are

SRL 6071  
(C-3021/2)

623

the same alkyl.

246. A compound of claim 234 wherein  $R^1$  and  $R^2$  are each n-butyl.

5

247. A compound of claim 234 wherein  
n is 1 or 2;

$R^1$  and  $R^2$  are n-butyl;

$R^3$  and  $R^6$  are hydrogen;

10


$R^4$  is hydroxy;

$R^7$  and  $R^8$  are hydrogen; and

one or more  $R^x$  are independently selected from methoxy and dimethylamino.


15

248. A compound of claim 234 having the structural formula:



249. A compound of formula (I):

20



(I)

wherein:

q is an integer from 1 to 4;

n is an integer from 0 to 2;

$R^1$  and  $R^2$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^wA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^w$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclylalkyl, and alkylammoniumalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which they are attached form  $C_3$ - $C_{10}$  cycloalkyl;

$R^3$  and  $R^4$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle,  $OR^9$ ,  $NR^9R^{10}$ ,  $SR^9$ ,  $S(O)R^9$ ,

SRL 6071  
(C-3021/2)

625

$\text{SO}_2\text{R}^9$ , and  $\text{SO}_3\text{R}^9$ , wherein  $\text{R}^9$  and  $\text{R}^{10}$  are as defined above; or

$\text{R}^3$  and  $\text{R}^4$  together form  $=\text{O}$ ,  $=\text{NOR}^{11}$ ,  $=\text{S}$ ,  $=\text{NNR}^{11}\text{R}^{12}$ ,  $=\text{NR}^9$ , or  $=\text{CR}^{11}\text{R}^{12}$ ,

5 wherein  $\text{R}^{11}$  and  $\text{R}^{12}$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
10  $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ ,  $\text{CO}_2\text{R}^9$ , CN, halogen, oxo, and  $\text{CONR}^9\text{R}^{10}$ , wherein  $\text{R}^9$  and  $\text{R}^{10}$  are as defined above, provided that both  $\text{R}^3$  and  $\text{R}^4$  cannot be OH,  $\text{NH}_2$ , and SH, or

$\text{R}^{11}$  and  $\text{R}^{12}$  together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

15  $\text{R}^5$  is aryl substituted with one or more substituent groups independently selected from the group consisting of  $\text{NR}^{13}\text{C}(\text{O})\text{R}^{14}$ ,  $\text{NR}^{13}\text{C}(\text{O})\text{NR}^{14}\text{R}^{15}$ ,  $\text{NR}^{13}\text{CO}_2\text{R}^{14}$ ,  $\text{OC}(\text{O})\text{R}^{13}$ ,  $\text{OC}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{NR}^{13}\text{SOR}^{14}$ ,  $\text{NR}^{13}\text{SO}_2\text{R}^{14}$ ,  $\text{NR}^{13}\text{SONR}^{14}\text{R}^{15}$ , and  $\text{NR}^{13}\text{SO}_2\text{NR}^{14}\text{R}^{15}$ ,

20 wherein:

$\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, arylalkyl, alkylarylalkyl, alkylheteroarylalkyl,  
25 alkylheterocyclalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary heterocycle, quaternary heteroaryl, heterocyclalkyl, heteroarylalkyl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, alkylammoniumalkyl, and  
30 carboxyalkylaminocarbonylalkyl,

$\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are optionally substituted with

SRL 6071  
(C-3021/2)

626

one or more groups selected from the group consisting of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl, heterocycle, heteroaryl, sulfoalkyl, quaternary heterocycle, quaternary heteroaryl, quaternary heterocyclylalkyl, quaternary heteroarylalkyl, guanidinyll,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S(O)}\text{R}^9$ ,  $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ , CN, halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO(OR}^{16})\text{OR}^{17}$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{R}^{11}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , and  $\text{C(O)OM}$ ,

wherein  $\text{A}^-$  is an pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation,

wherein  $\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected from the substituents constituting  $\text{R}^9$  and M; or

$\text{R}^{13}$  and  $\text{R}^{14}$ , together with the nitrogen atom to which they are attached form a mono- or polycyclic heterocycle that is optionally substituted with one or more radicals selected from the group consisting of oxo, carboxy and quaternary salts; or

$\text{R}^{14}$  and  $\text{R}^{15}$ , together with the nitrogen atom to which they are attached, form a cyclic ring; and

$\text{R}^6$  is selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle,  $\text{OR}^{30}$ ,  $\text{SR}^9$ ,  $\text{S(O)}\text{R}^9$ ,  $\text{SO}_2\text{R}^9$ , and  $\text{SO}_3\text{R}^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary



heteroaryl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  
 $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  
 OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  
 $NR^{13}C(O)R^{14}$ ,  $NR^{13}C(O)NR^{14}R^{15}$ ,  $NR^{13}CO_2R^{14}$ ,  $OC(O)R^{13}$ ,  
 5  $OC(O)NR^{13}R^{14}$ ,  $NR^{13}SOR^{14}$ ,  $NR^{13}SO_2R^{14}$ ,  $NR^{13}SONR^{14}R^{15}$ ,  
 $NR^{13}SO_2NR^{14}R^{15}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  
 $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,

wherein:

10  $A^-$  is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $OR^7$ ,  
 15  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

20 wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,  $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  $P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are  
 25 independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, arylalkyl, alkylarylalkyl, alkylheteroarylalkyl, alkylheterocyclalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary  
 30 heterocycle, quaternary heteroaryl, heterocyclalkyl,

heteroarylalkyl, quaternary heterocyclalkyl,  
quaternary heteroarylalkyl, alkylammoniumalkyl, and  
carboxyalkylaminocarbonylalkyl,

5 wherein alkyl, alkenyl, alkynyl, arylalkyl,  
heterocycle, and polyalkyl optionally have one or more  
carbons replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  
 $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ ,  $\text{P}(\text{O})\text{R}^9$ , phenylene, carbohydrate,  
amino acid, peptide, or polypeptide, and

10  $\text{R}^{13}$ ,  $\text{R}^{14}$ , and  $\text{R}^{15}$  are optionally substituted with  
one or more groups selected from the group consisting  
of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl,  
heterocycle, heteroaryl, sulfoalkyl, quaternary  
heterocycle, quaternary heteroaryl, quaternary  
heterocyclalkyl, quaternary heteroarylalkyl,  
15 guanidinyll,  $\text{OR}^9$ ,  $\text{NR}^9\text{R}^{10}$ ,  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ ,  
 $\text{SO}_2\text{R}^9$ ,  $\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ , CN, halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_2\text{OM}$ ,  
 $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{R}^{11}\text{A}^-$ ,  $\text{S}^+\text{R}^9\text{R}^{10}\text{A}^-$ , and  
 $\text{C}(\text{O})\text{OM}$ ,

20 wherein  $\text{R}^{16}$  and  $\text{R}^{17}$  are independently selected  
from the substituents constituting  $\text{R}^9$  and M; or

$\text{R}^{13}$  and  $\text{R}^{14}$ , together with the nitrogen atom to  
which they are attached form a mono- or polycyclic  
heterocycle that is optionally substituted with one or  
more radicals selected from the group consisting of  
25 oxo, carboxy and quaternary salts; or

$\text{R}^{14}$  and  $\text{R}^{15}$ , together with the nitrogen atom to  
which they are attached, form a cyclic ring; and

30  $\text{R}^{30}$  is selected from the group consisting of alkyl,  
alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
ammoniumalkyl, alkylammoniumalkyl, arylalkyl,  
carboxyalkyl, carboxyheteroaryl, carboxyheterocycle,  
carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,

heterocyclalkyl, and alkylammoniumalkyl; and

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl; and

one or more  $R^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,

SRL 6071  
(C-3021/2)

630

SO<sub>2</sub>OM, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, PO(OR<sup>16</sup>)OR<sup>17</sup>, and C(O)OM,

wherein in R<sup>x</sup>, one or more carbons are optionally replaced by O, NR<sup>13</sup>, N<sup>+</sup>R<sup>13</sup>R<sup>14</sup>A<sup>-</sup>, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>13</sup>A<sup>-</sup>, PR<sup>13</sup>, P(O)R<sup>13</sup>, P<sup>+</sup>R<sup>13</sup>R<sup>14</sup>A<sup>-</sup>, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR<sup>9</sup>, N<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>9</sup>A<sup>-</sup>, PR<sup>9</sup>, P<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, or P(O)R<sup>9</sup>;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR<sup>13</sup>, NR<sup>13</sup>R<sup>14</sup>, SR<sup>13</sup>, S(O)R<sup>13</sup>, SO<sub>2</sub>R<sup>13</sup>, SO<sub>3</sub>R<sup>13</sup>, NR<sup>13</sup>OR<sup>14</sup>, NR<sup>13</sup>NR<sup>14</sup>R<sup>15</sup>, NO<sub>2</sub>, CO<sub>2</sub>R<sup>13</sup>, CN, OM, SO<sub>2</sub>OM, SO<sub>2</sub>NR<sup>13</sup>R<sup>14</sup>, C(O)NR<sup>13</sup>R<sup>14</sup>, C(O)OM, COR<sup>13</sup>, P(O)R<sup>13</sup>R<sup>14</sup>, P<sup>+</sup>R<sup>13</sup>R<sup>14</sup>R<sup>15</sup>A<sup>-</sup>, P(OR<sup>13</sup>)OR<sup>14</sup>, S<sup>+</sup>R<sup>13</sup>R<sup>14</sup>A<sup>-</sup>, and N<sup>+</sup>R<sup>9</sup>R<sup>11</sup>R<sup>12</sup>A<sup>-</sup>, or

a pharmaceutically acceptable salt, solvate, or prodrug thereof.

250. A compound of claim 249 wherein R<sup>5</sup> is aryl substituted with a radical selected from the group consisting of NR<sup>13</sup>C(O)NR<sup>14</sup>R<sup>15</sup> and NR<sup>13</sup>CO<sub>2</sub>R<sup>14</sup>.

251. A compound of claim 249 wherein R<sup>5</sup> is phenyl substituted with a radical selected from the group consisting of NR<sup>13</sup>C(O)NR<sup>14</sup>R<sup>15</sup> and NR<sup>13</sup>CO<sub>2</sub>R<sup>14</sup>.

252. A compound of claim 249 wherein n is 1 or 2.

5 253. A compound of claim 249 wherein  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen and alkyl.

10 254. A compound of claim 249 wherein  $R^7$  and  $R^8$  are hydrogen.

255. A compound of claim 249 wherein  $R^3$  and  $R^4$  are independently selected from the group consisting of hydrogen and  $OR^9$ .

15 256. A compound of claim 249 wherein  $R^3$  is hydrogen and  $R^4$  is hydroxy.

20 257. A compound of claim 249 wherein one or more  $R^x$  are independently selected from the group consisting of  $OR^{13}$  and  $NR^{13}R^{14}$ .

25 258. A compound of claim 249 wherein one or more  $R^x$  are independently selected from methoxy and dimethylamino.

259. A compound of claim 249 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen and alkyl.

30 260. A compound of claim 249 wherein  $R^1$  and  $R^2$  are independently selected from the group consisting alkyl.

35 261. A compound of claim 249 wherein  $R^1$  and  $R^2$  are the same alkyl.

SRL 6071  
(C-3021/2)

632

262. A compound of claim 249 wherein  $R^1$  and  $R^2$  are each n-butyl.

263. A compound of claim 249 wherein  
n is 1 or 2;

$R^1$  and  $R^2$  are n-butyl;


$R^3$  and  $R^6$  are hydrogen;

$R^4$  is hydroxy;

$R^7$  and  $R^8$  are hydrogen; and

one or more  $R^x$  are independently selected from methoxy and dimethylamino.

264. A compound of claim 98 having the structural formula:



265. A pharmaceutical composition comprising an anti-hyperlipidemic condition effective amount of a compound of formula (I) of claim 170, and

SRL 6071  
(C-3021/2)

633

a pharmaceutically acceptable carrier.

5        266. A pharmaceutical composition comprising an  
anti-atherosclerotic effective amount of a compound of  
formula (I) of claim 170, and  
a pharmaceutically acceptable carrier.

10       267. A pharmaceutical composition comprising an  
anti-hypercholesterolemia effective amount of a  
compound of formula (I) of claim 170, and  
a pharmaceutically acceptable carrier.

15       268. A method for the prophylaxis or treatment of  
a hyperlipidemic condition comprising administering to  
a patient in need thereof a composition of claim 265 in  
unit dosage form.

20       269. A method for the prophylaxis or treatment of  
an atherosclerotic condition comprising administering  
to a patient in need thereof a composition of claim 266  
in unit dosage form.

25       270. A method for the prophylaxis or treatment of  
hypercholesterolemia comprising administering to a  
patient in need thereof a composition of claim 267 in  
unit dosage form.

30       271. A pharmaceutical composition comprising an  
anti-hyperlipidemic condition effective amount of a  
compound of formula (I) of claim 180, and  
a pharmaceutically acceptable carrier.

35       272. A pharmaceutical composition comprising an  
anti-atherosclerotic effective amount of a compound of  
formula (I) of claim 180, and

a pharmaceutically acceptable carrier.

5        273. A pharmaceutical composition comprising an  
anti-hypercholesterolemia effective amount of a  
compound of formula (I) of claim 180, and  
a pharmaceutically acceptable carrier.

10       274. A method for the prophylaxis or treatment of  
a hyperlipidemic condition comprising administering to  
a patient in need thereof a composition of claim 271 in  
unit dosage form.

15       275. A method for the prophylaxis or treatment of  
an atherosclerotic condition comprising administering  
to a patient in need thereof a composition of claim 272  
in unit dosage form.

20       276. A method for the prophylaxis or treatment of  
hypercholesterolemia comprising administering to a  
patient in need thereof a composition of claim 273 in  
unit dosage form.

25       277. A pharmaceutical composition comprising an  
anti-hyperlipidemic condition effective amount of a  
compound of formula (I) of claim 201, and  
a pharmaceutically acceptable carrier.

30       278. A pharmaceutical composition comprising an  
anti-atherosclerotic effective amount of a compound of  
formula (I) of claim 201, and  
a pharmaceutically acceptable carrier.

35       279. A pharmaceutical composition comprising an  
anti-hypercholesterolemia effective amount of a  
compound of formula (I) of claim 201, and



a pharmaceutically acceptable carrier.

5 280. A method for the prophylaxis or treatment of  
a hyperlipidemic condition comprising administering to  
a patient in need thereof a composition of claim 277 in  
unit dosage form.

10 281. A method for the prophylaxis or treatment of  
an atherosclerotic condition comprising administering  
to a patient in need thereof a composition of claim 278  
in unit dosage form.

15 282. A method for the prophylaxis or treatment of  
hypercholesterolemia comprising administering to a  
patient in need thereof a composition of claim 279 in  
unit dosage form.

20 283. A pharmaceutical composition comprising an  
anti-hyperlipidemic condition effective amount of a  
compound of formula (I) of claim 218, and  
a pharmaceutically acceptable carrier.

25 284. A pharmaceutical composition comprising an  
anti-atherosclerotic effective amount of a compound of  
formula (I) of claim 218, and  
a pharmaceutically acceptable carrier.

30 285. A pharmaceutical composition comprising an  
anti-hypercholesterolemia effective amount of a  
compound of formula (I) of claim 218, and  
a pharmaceutically acceptable carrier.

35 286. A method for the prophylaxis or treatment of  
a hyperlipidemic condition comprising administering to  
a patient in need thereof a composition of claim 283 in

unit dosage form.

287. A method for the prophylaxis or treatment of  
an atherosclerotic condition comprising administering  
5 to a patient in need thereof a composition of claim 284  
in unit dosage form.

288. A method for the prophylaxis or treatment of  
hypercholesterolemia comprising administering to a  
10 patient in need thereof a composition of claim 285 in  
unit dosage form.

289. A pharmaceutical composition comprising an  
anti-hyperlipidemic condition effective amount of a  
15 compound of formula (I) of claim 234, and  
a pharmaceutically acceptable carrier.

290. A pharmaceutical composition comprising an  
anti-atherosclerotic effective amount of a compound of  
20 formula (I) of claim 234, and  
a pharmaceutically acceptable carrier.

291. A pharmaceutical composition comprising an  
anti-hypercholesterolemia effective amount of a  
25 compound of formula (I) of claim 234, and  
a pharmaceutically acceptable carrier.

292. A method for the prophylaxis or treatment of  
a hyperlipidemic condition comprising administering to  
30 a patient in need thereof a composition of claim 289 in  
unit dosage form.

293. A method for the prophylaxis or treatment of  
an atherosclerotic condition comprising administering  
35 to a patient in need thereof a composition of claim 290

SRL 6071  
(C-3021/2)

637

in unit dosage form.

5           294. A method for the prophylaxis or treatment of hypercholesterolemia comprising administering to a patient in need thereof a composition of claim 291 in unit dosage form.

10           295. A pharmaceutical composition comprising an anti-hyperlipidemic condition effective amount of a compound of formula (I) of claim 249, and a pharmaceutically acceptable carrier.

15           296. A pharmaceutical composition comprising an anti-atherosclerotic effective amount of a compound of formula (I) of claim 249, and a pharmaceutically acceptable carrier.

20           297. A pharmaceutical composition comprising an anti-hypercholesterolemia effective amount of a compound of formula (I) of claim 249, and a pharmaceutically acceptable carrier.


25           298. A method for the prophylaxis or treatment of a hyperlipidemic condition comprising administering to a patient in need thereof a composition of claim 295 in unit dosage form.

30           299. A method for the prophylaxis or treatment of an atherosclerotic condition comprising administering to a patient in need thereof a composition of claim 296 in unit dosage form.

35           300. A method for the prophylaxis or treatment of hypercholesterolemia comprising administering to a patient in need thereof a composition of claim 297 in

unit dosage form.

301. A process for the preparation of a compound having the formula:



XLI

comprising:

treating a thiophenol with an abstracting agent;  
coupling the thiophenyl and a cyclic sulfate to  
form an intermediate comprising a sulfate group; and  
removing the sulfate group of the intermediate to  
form the compound of formula XLI;

wherein

$q$  is an integer from 1 to 4;

$R^1$  and  $R^2$  are independently selected from the group  
consisting of H, alkyl, alkenyl, alkynyl, haloalkyl,  
alkylaryl, arylalkyl, alkoxy, alkoxyalkyl,  
dialkylamino, alkylthio, (polyalkyl)aryl, and  
cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl,  
alkylaryl, arylalkyl, alkoxy, alkoxyalkyl,  
dialkylamino, alkylthio, (polyalkyl)aryl, and  
cycloalkyl optionally are substituted with one or more  
substituents selected from the group consisting of  $OR^9$ ,  
 $NR^9R^{10}$ ,  $N^+R^9R^{10}R^wA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  
 $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl,  
alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl  
optionally have one or more carbons replaced by O,  $NR^9$ ,

$N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,  
wherein R<sup>9</sup>, R<sup>10</sup>, and R<sup>v</sup> are independently selected  
from the group consisting of H, alkyl, alkenyl,  
alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
5 ammoniumalkyl, arylalkyl, carboxyalkyl,  
carboxyheteroaryl, carboxyheterocycle,  
carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,  
heterocyclylalkyl, and alkylammoniumalkyl; or

R<sup>1</sup> and R<sup>2</sup> taken together with the carbon to which  
10 they are attached form C<sub>3</sub>-C<sub>10</sub> cycloalkyl;

R<sup>3</sup> is hydroxy;

R<sup>4</sup> is hydrogen;

R<sup>5</sup> and R<sup>6</sup> are independently selected from the  
group consisting of H, alkyl, alkenyl, alkynyl, aryl,  
15 cycloalkyl, heterocycle, quaternary heterocycle, OR<sup>30</sup>,  
SR<sup>9</sup>, S(O)R<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup>, and SO<sub>3</sub>R<sup>9</sup>,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl,  
heterocycle, quaternary heterocycle, and quaternary  
heteroaryl can be substituted with one or more  
20 substituent groups independently selected from the  
group consisting of alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, heterocycle,  
arylalkyl, quaternary heterocycle, quaternary  
heteroaryl, halogen, oxo, OR<sup>13</sup>, NR<sup>13</sup>R<sup>14</sup>, SR<sup>13</sup>, S(O)R<sup>13</sup>,  
25 SO<sub>2</sub>R<sup>13</sup>, SO<sub>3</sub>R<sup>13</sup>, NR<sup>13</sup>OR<sup>14</sup>, NR<sup>13</sup>NR<sup>14</sup>R<sup>15</sup>, NO<sub>2</sub>, CO<sub>2</sub>R<sup>13</sup>, CN,  
OM, SO<sub>2</sub>OM, SO<sub>2</sub>NR<sup>13</sup>R<sup>14</sup>, C(O)NR<sup>13</sup>R<sup>14</sup>, C(O)OM, COR<sup>13</sup>,  
NR<sup>13</sup>C(O)R<sup>14</sup>, NR<sup>13</sup>C(O)NR<sup>14</sup>R<sup>15</sup>, NR<sup>13</sup>CO<sub>2</sub>R<sup>14</sup>, OC(O)R<sup>13</sup>,  
OC(O)NR<sup>13</sup>R<sup>14</sup>, NR<sup>13</sup>SOR<sup>14</sup>, NR<sup>13</sup>SO<sub>2</sub>R<sup>14</sup>, NR<sup>13</sup>SONR<sup>14</sup>R<sup>15</sup>,  
NR<sup>13</sup>SO<sub>2</sub>NR<sup>14</sup>R<sup>15</sup>, P(O)R<sup>13</sup>R<sup>14</sup>, P<sup>+</sup>R<sup>13</sup>R<sup>14</sup>R<sup>15</sup>A<sup>-</sup>, P(OR<sup>13</sup>)OR<sup>14</sup>,  
30 S<sup>+</sup>R<sup>13</sup>R<sup>14</sup>A<sup>-</sup>, and N<sup>+</sup>R<sup>9</sup>R<sup>11</sup>R<sup>12</sup>A<sup>-</sup>,

wherein:

A<sup>-</sup> is a pharmaceutically acceptable anion and M is

a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent

5 groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

10 wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,  $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,  $P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are  
15 independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, arylalkyl, alkylarylalkyl, alkylheteroarylalkyl, alkylheterocyclalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary  
20 heterocycle, quaternary heteroaryl, heterocyclalkyl, heteroarylalkyl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, alkylammoniumalkyl, and carboxyalkylaminocarbonylalkyl,

25 wherein alkyl, alkenyl, alkynyl, arylalkyl, heterocycle, and polyalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ ,  $P(O)R^9$ , phenylene, carbohydrate, amino acid, peptide, or polypeptide, and

30  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are optionally substituted with one or more groups selected from the group consisting

of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl,  
heterocycle, heteroaryl, sulfoalkyl, quaternary  
heterocycle, quaternary heteroaryl, quaternary  
heterocyclylalkyl, quaternary heteroarylalkyl,  
5 guanidiny,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  
 $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  
 $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S^+R^9R^{10}A^-$ , and  
C(O)OM,

wherein  $R^{16}$  and  $R^{17}$  are independently selected  
10 from the substituents constituting  $R^9$  and M; or  
 $R^{13}$  and  $R^{14}$ , together with the nitrogen atom to  
which they are attached form a mono- or polycyclic  
heterocycle that is optionally substituted with one or  
more radicals selected from the group consisting of  
15 oxo, carboxy and quaternary salts; or

$R^{14}$  and  $R^{15}$ , together with the nitrogen atom to  
which they are attached, form a cyclic ring; and  
 $R^{30}$  is selected from the group consisting of alkyl,  
alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
20 ammoniumalkyl, alkylammoniumalkyl, arylalkyl,  
carboxyalkyl, carboxyheteroaryl, carboxyheterocycle,  
carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,  
heterocyclylalkyl, and alkylammoniumalkyl; and

$R^7$  and  $R^8$  are hydrogen; and  
25 one or more  $R^*$  are independently selected from the  
group consisting of H, alkyl, alkenyl, alkynyl,  
polyalkyl, acyloxy, aryl, arylalkyl, halogen,  
haloalkyl, cycloalkyl, heterocycle, heteroaryl,  
polyether, quaternary heterocycle, quaternary  
30 heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  
 $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  
 $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  
 $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,

SRL 6071  
(C-3021/2)

642

$P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, and alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

wherein in  $R^x$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,


wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ ;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,



$P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ .


302. The process of claim 301 wherein the cyclic sulfate has the formula:



5

XL

and the thiophenol has the formula:



XVIII A

wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>\*</sup> and q are as defined in claim 301.

10

303. The process of claim 301 wherein the sulfate group is removed by treating the intermediate with a hydrolyzing agent.

15

304. The process of claim 303 wherein the hydrolyzing agent is a mineral acid.

305. The process of claim 303 wherein the

hydrolyzing agent is selected from the group consisting of hydrochloric acid and sulfuric acid.

5 306. The process of claim 302 wherein the abstracting agent is a base having a pH of at least about 10.

10 307. The process of claim 302 wherein the abstracting agent is an alkali metal hydride.


308. The process of claim 302 wherein the abstracting agent is sodium hydride.

15 309. The process of claim 302 wherein the  $R^1$  and  $R^2$  are alkyl.

20 310. The process of claim 302 wherein the  $R^1$  and  $R^2$  are selected from the group consisting of ethyl, n-butyl, iso-butyl and pentyl.

311. The process of claim 302 wherein the  $R^1$  and  $R^2$  are n-butyl.

25 312. A process for the preparation of a compound having the formula I:



(I)

comprising:

reacting a cyclic sulfate with a thiophenol to  
form an alcohol;

oxidizing said alcohol to form a sulfone-aldehyde;  
and

5 cyclizing said sulfone-aldehyde to form the  
compound of formula I;

wherein:

q is an integer from 1 to 4;

n is 2;

10  $R^1$  and  $R^2$  are independently selected from the group  
consisting of H, alkyl, alkenyl, alkynyl, haloalkyl,  
alkylaryl, arylalkyl, alkoxy, alkoxyalkyl,  
dialkylamino, alkylthio, (polyalkyl)aryl, and  
cycloalkyl,

15 wherein alkyl, alkenyl, alkynyl, haloalkyl,  
alkylaryl, arylalkyl, alkoxy, alkoxyalkyl,  
dialkylamino, alkylthio, (polyalkyl)aryl, and  
cycloalkyl optionally are substituted with one or more  
substituents selected from the group consisting of  $OR^9$ ,  
20  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^wA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  
 $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

wherein alkyl, alkenyl, alkynyl, alkylaryl,  
alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl  
optionally have one or more carbons replaced by O,  $NR^9$ ,  
25  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

wherein  $R^9$ ,  $R^{10}$ , and  $R^w$  are independently selected  
from the group consisting of H, alkyl, alkenyl,  
alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
ammoniumalkyl, arylalkyl, carboxyalkyl,  
30 carboxyheteroaryl, carboxyheterocycle,  
carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,  
heterocyclylalkyl, and alkylammoniumalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which  
they are attached form  $C_3$ - $C_{10}$  cycloalkyl;

35  $R^3$  is hydroxy;

$R^4$  is hydrogen;

$R^5$  and  $R^6$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle,  $OR^{30}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $NR^{13}C(O)R^{14}$ ,  $NR^{13}C(O)NR^{14}R^{15}$ ,  $NR^{13}CO_2R^{14}$ ,  $OC(O)R^{13}$ ,  $OC(O)NR^{13}R^{14}$ ,  $NR^{13}SOR^{14}$ ,  $NR^{13}SO_2R^{14}$ ,  $NR^{13}SONR^{14}R^{15}$ ,  $NR^{13}SO_2NR^{14}R^{15}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,

wherein:

$A^-$  is a pharmaceutically acceptable anion and  $M$  is a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,

$P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,

5  $NR^7$ ,  $N^+R^7R^8A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,

$P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl,

polyether, aryl, arylalkyl, alkylarylalkyl, 10 alkylheteroarylalkyl, alkylheterocyclalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary heterocycle, quaternary heteroaryl, heterocyclalkyl, heteroarylalkyl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, alkylammoniumalkyl, and 15 carboxyalkylaminocarbonylalkyl,

wherein alkyl, alkenyl, alkynyl, arylalkyl, heterocycle, and polyalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ ,  $P(O)R^9$ , phenylene, carbohydrate, 20 amino acid, peptide, or polypeptide, and

$R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are optionally substituted with one or more groups selected from the group consisting of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl, heterocycle, heteroaryl, sulfoalkyl, quaternary 25 heterocycle, quaternary heteroaryl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, guanidiny,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S^+R^9R^{10}A^-$ , and 30  $C(O)OM$ ,

wherein  $R^{16}$  and  $R^{17}$  are independently selected

from the substituents constituting  $R^9$  and M; or  
 $R^{13}$  and  $R^{14}$ , together with the nitrogen atom to  
 which they are attached form a mono- or polycyclic  
 heterocycle that is optionally substituted with one or  
 5 more radicals selected from the group consisting of  
 oxo, carboxy and quaternary salts; or

$R^{14}$  and  $R^{15}$ , together with the nitrogen atom to  
 which they are attached, form a cyclic ring; and

$R^{30}$  is selected from the group consisting of alkyl,  
 10 alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
 ammoniumalkyl, alkylammoniumalkyl, arylalkyl,  
 carboxyalkyl, carboxyheteroaryl, carboxyheterocycle,  
 carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,  
 heterocyclylalkyl, and alkylammoniumalkyl; and

15  $R^7$  and  $R^8$  are hydrogen; and

one or more  $R^*$  are independently selected from the  
 group consisting of H, alkyl, alkenyl, alkynyl,  
 polyalkyl, acyloxy, aryl, arylalkyl, halogen,  
 haloalkyl, cycloalkyl, heterocycle, heteroaryl,  
 20 polyether, quaternary heterocycle, quaternary  
 heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  
 $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  
 $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  
 $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  
 25  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and  
 carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl,  
 polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl,  
 polyether, quaternary heterocycle, and quaternary  
 30 heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  
 $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN,  
 halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  
 $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting

of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle,  
heteroaryl, and alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl,  
heterocycle, heteroaryl, alkyl, quaternary heterocycle,  
5 and quaternary heteroaryl optionally are substituted  
with one or more substituents selected from the group  
consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  
 $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2R^9$ ,  $SO_2OM$ ,  
 $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

10 wherein in  $R^x$ , one or more carbons are optionally  
replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  
 $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide,  
polypeptide, carbohydrate, polyether, or polyalkyl,


wherein in said polyalkyl, phenylene, amino acid,  
15 peptide, polypeptide, and carbohydrate, one or more  
carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S,  
SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ ;

wherein quaternary heterocycle and quaternary  
heteroaryl are optionally substituted with one or more  
20 groups selected from the group consisting of alkyl,  
alkenyl, alkynyl, polyalkyl, polyether, aryl,  
haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,  
oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  
 $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  
25  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  
 $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ .

313. The process of claim 312 wherein the cyclic  
sulfate has the formula:


SRL 6071  
(C-3021/2)

650



XL

and the thiophenol has the formula:



XVIII A

wherein  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^x$  and  $q$  are as defined in claim 312.

314. The process of claim 313 wherein the  $R^1$  and  $R^2$  are alkyl.

315. The process of claim 313 wherein the  $R^1$  and  $R^2$  are selected from the group consisting of ethyl, n-butyl, iso-butyl and pentyl.

316. The process of claim 313 wherein the  $R^1$  and  $R^2$  are n-butyl.

317. The process of claim 313 wherein the alcohol is oxidized with an oxidizing agent to form an



SRL 6071  
(C-3021/2)

651

aldehyde.

318. The process of claim 317 wherein the aldehyde is oxidized with an oxidizing agent to form a sulfone-aldehyde.


319. The process of claim 313 wherein the sulfone-aldehyde is cyclized with a cyclizing agent that is a base having a pH between about 8 to about 9.

320. The process of claim 313 wherein the sulfone-aldehyde is cyclized with a cyclizing agent that is an alkali alkoxide base.

321. The process of claim 313 wherein the sulfone-aldehyde is cyclized with potassium tert-butoxide.

322. The process of claim 313 wherein the alcohol is oxidized with pyridinium chlorochromate to form an aldehyde; the aldehyde is oxidized with metachloroperbenzoic acid to form a sulfone-aldehyde; and the sulfone-aldehyde is cyclized with potassium tert-butoxide.

323. A process for the preparation of a compound having the formula LI:



comprising:

treating a halobenzene with an abstracting agent;

SRL 6071  
(C-3021/2)

652

coupling the halobenzene and a cyclic sulfate to  
form an intermediate comprising a sulfate group; and  
removing the sulfate group of the intermediate to  
form the compound of formula LI; wherein

5        q is an integer from 1 to 4;

$R^1$  and  $R^2$  are independently selected from the group  
consisting of H, alkyl, alkenyl, alkynyl, haloalkyl,  
alkylaryl, arylalkyl, alkoxy, alkoxyalkyl,  
dialkylamino, alkylthio, (polyalkyl)aryl, and  
10        cycloalkyl,

      wherein alkyl, alkenyl, alkynyl, haloalkyl,  
alkylaryl, arylalkyl, alkoxy, alkoxyalkyl,  
dialkylamino, alkylthio, (polyalkyl)aryl, and  
cycloalkyl optionally are substituted with one or more  
15        substituents selected from the group consisting of  $OR^9$ ,  
 $NR^9R^{10}$ ,  $N^+R^9R^{10}R^wA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  
 $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

      wherein alkyl, alkenyl, alkynyl, alkylaryl,  
alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl  
20        optionally have one or more carbons replaced by O,  $NR^9$ ,  
 $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,  
wherein  $R^9$ ,  $R^{10}$ , and  $R^w$  are independently selected

from the group consisting of H, alkyl, alkenyl,  
alkynyl, cycloalkyl, aryl, acyl, heterocycle,  
25        ammoniumalkyl, arylalkyl, carboxyalkyl,  
carboxyheteroaryl, carboxyheterocycle,  
carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl,  
heterocyclylalkyl, and alkylammoniumalkyl; or

$R^1$  and  $R^2$  taken together with the carbon to which  
30        they are attached form  $C_3$ - $C_{10}$  cycloalkyl;

$R^3$  is hydroxy;

$R^4$  is hydrogen;

$R^5$  and  $R^6$  are independently selected from the  
group consisting of H, alkyl, alkenyl, alkynyl, aryl,

cycloalkyl, heterocycle, quaternary heterocycle,  $OR^{30}$ ,  
 $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl,  
heterocycle, quaternary heterocycle, and quaternary  
heteroaryl can be substituted with one or more  
substituent groups independently selected from the  
group consisting of alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, heterocycle,  
arylalkyl, quaternary heterocycle, quaternary  
heteroaryl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  
 $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  
 $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  
 $NR^{13}C(O)R^{14}$ ,  $NR^{13}C(O)NR^{14}R^{15}$ ,  $NR^{13}CO_2R^{14}$ ,  $OC(O)R^{13}$ ,  
 $OC(O)NR^{13}R^{14}$ ,  $NR^{13}SOR^{14}$ ,  $NR^{13}SO_2R^{14}$ ,  $NR^{13}SONR^{14}R^{15}$ ,  
 $NR^{13}SO_2NR^{14}R^{15}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  
 $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,

wherein:

$A^-$  is a pharmaceutically acceptable anion and  $M$  is  
a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, and heterocycle  
can be further substituted with one or more substituent  
groups selected from the group consisting of  $OR^7$ ,  
 $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  
 $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl,  
cycloalkyl, heterocycle, arylalkyl, quaternary  
heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  
 $P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, haloalkyl, cycloalkyl, and heterocycle

can optionally have one or more carbons replaced by O,  
NR<sup>7</sup>, N<sup>+</sup>R<sup>7</sup>R<sup>8</sup>A<sup>-</sup>, S, SO, SO<sub>2</sub>, S<sup>+</sup>R<sup>7</sup>A<sup>-</sup>, PR<sup>7</sup>, P(O)R<sup>7</sup>,  
P<sup>+</sup>R<sup>7</sup>R<sup>8</sup>A<sup>-</sup>, or phenylene, and R<sup>13</sup>, R<sup>14</sup>, and R<sup>15</sup> are  
independently selected from the group consisting of  
5 hydrogen, alkyl, alkenyl, alkynyl, polyalkyl,  
polyether, aryl, arylalkyl, alkylarylalkyl,  
alkylheteroarylalkyl, alkylheterocyclalkyl,  
cycloalkyl, heterocycle, heteroaryl, quaternary  
heterocycle, quaternary heteroaryl, heterocyclalkyl,  
10 heteroarylalkyl, quaternary heterocyclalkyl,  
quaternary heteroarylalkyl, alkylammoniumalkyl, and  
carboxyalkylaminocarbonylalkyl,

wherein alkyl, alkenyl, alkynyl, arylalkyl,  
heterocycle, and polyalkyl optionally have one or more  
15 carbons replaced by O, NR<sup>9</sup>, N<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, S, SO, SO<sub>2</sub>,  
S<sup>+</sup>R<sup>9</sup>A<sup>-</sup>, PR<sup>9</sup>, P<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, P(O)R<sup>9</sup>, phenylene, carbohydrate,  
amino acid, peptide, or polypeptide, and

R<sup>13</sup>, R<sup>14</sup>, and R<sup>15</sup> are optionally substituted with  
one or more groups selected from the group consisting  
20 of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl,  
heterocycle, heteroaryl, sulfoalkyl, quaternary  
heterocycle, quaternary heteroaryl, quaternary  
heterocyclalkyl, quaternary heteroarylalkyl,  
guanidiny, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, N<sup>+</sup>R<sup>9</sup>R<sup>11</sup>R<sup>12</sup>A<sup>-</sup>, SR<sup>9</sup>, S(O)R<sup>9</sup>,  
25 SO<sub>2</sub>R<sup>9</sup>, SO<sub>3</sub>R<sup>9</sup>, oxo, CO<sub>2</sub>R<sup>9</sup>, CN, halogen, CONR<sup>9</sup>R<sup>10</sup>, SO<sub>2</sub>OM,  
SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, PO(OR<sup>16</sup>)OR<sup>17</sup>, P<sup>+</sup>R<sup>9</sup>R<sup>10</sup>R<sup>11</sup>A<sup>-</sup>, S<sup>+</sup>R<sup>9</sup>R<sup>10</sup>A<sup>-</sup>, and  
C(O)OM,

wherein R<sup>16</sup> and R<sup>17</sup> are independently selected  
from the substituents constituting R<sup>9</sup> and M; or  
30 R<sup>13</sup> and R<sup>14</sup>, together with the nitrogen atom to  
which they are attached form a mono- or polycyclic

heterocycle that is optionally substituted with one or more radicals selected from the group consisting of oxo, carboxy and quaternary salts; or

5  $R^{14}$  and  $R^{15}$ , together with the nitrogen atom to which they are attached, form a cyclic ring; and

$R^{30}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclalkyl, and alkylammoniumalkyl; and

$R^7$  and  $R^8$  are hydrogen; and

one or more  $R^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

25 wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

30 wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, and alkyl,

35 wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle,

and quaternary heteroaryl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_3R^9$ ,  $SO_2OM$ ,  
5  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ , and  $C(O)OM$ ,

wherein in  $R^x$ , one or more carbons are optionally replaced by O,  $NR^{13}$ ,  $N^+R^{13}R^{14}A^-$ , S, SO,  $SO_2$ ,  $S^+R^{13}A^-$ ,  $PR^{13}$ ,  $P(O)R^{13}$ ,  $P^+R^{13}R^{14}A^-$ , phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

10 wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ , or  $P(O)R^9$ ;


wherein quaternary heterocycle and quaternary  
15 heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  
20  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ , CN, OM,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ; and

$R^e$  is an electron-withdrawing group located at the para or ortho position.

25 324. The process of claim 323 wherein the cyclic sulfate has the formula:


SRL 6071  
(C-3021/2)

657



XL

and the halobenzene has the formula:



wherein  $R^h$  is halogen, and  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^x$ ,  $R^e$  and  $q$   
are as defined in claim 323.

325. The process of claim 324 wherein the sulfate group is removed by treating the intermediate with a hydrolyzing agent.

326. The process of claim 325 wherein the hydrolyzing agent is a mineral acid.

327. The process of claim 325 wherein the hydrolyzing agent is selected from the group consisting of hydrochloric acid and sulfuric acid.

328. The process of claim 324 wherein the abstracting agent is a dialkali metal sulfide.

329. The process of claim 324 wherein the abstracting agent is dilithium sulfide.

330. The process of claim 324 wherein  $R^1$  and  $R^2$  are alkyl.


5 331. The process of claim 324 wherein  $R^1$  and  $R^2$  are selected from the group consisting of ethyl, n-butyl, iso-butyl and pentyl.

10 332. The process of claim 324 wherein  $R^1$  and  $R^2$  are n-butyl.

333. The process of claim 324 wherein  $R^h$  is chloro.

15 334. The process of claim 324 wherein  $R^e$  is p-nitro.

335. A process for the preparation of a compound having the formula I:



20 comprising:  
reacting a cyclic sulfate with a halobenzene to  
form an alcohol;  
oxidizing said alcohol to form a sulfone-aldehyde;  
25 and  
cyclizing said sulfone-aldehyde to form the  
compound of formula I;



wherein

q is an integer from 1 to 4;

n is 2;

5  $R^1$  and  $R^2$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

10 wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{10}R^wA^-$ ,  $SR^9$ ,  $S^+R^9R^{10}A^-$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ ,  $CO_2R^9$ , CN, halogen, oxo, and  $CONR^9R^{10}$ ,

15 wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, (polyalkyl)aryl, and cycloalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO,  $SO_2$ ,  $S^+R^9A^-$ ,  $P^+R^9R^{10}A^-$ , or phenylene,

20 wherein  $R^9$ ,  $R^{10}$ , and  $R^w$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclalkyl, and alkylammoniumalkyl; or

25  $R^1$  and  $R^2$  taken together with the carbon to which they are attached form  $C_3$ - $C_{10}$  cycloalkyl;

$R^3$  is hydroxy;

30  $R^4$  is hydrogen;

$R^5$  and  $R^6$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle,  $OR^{30}$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ , and  $SO_3R^9$ ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, halogen, oxo,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $SO_2R^{13}$ ,  $SO_3R^{13}$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $C(O)NR^{13}R^{14}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $NR^{13}C(O)R^{14}$ ,  $NR^{13}C(O)NR^{14}R^{15}$ ,  $NR^{13}CO_2R^{14}$ ,  $OC(O)R^{13}$ ,  $OC(O)NR^{13}R^{14}$ ,  $NR^{13}SOR^{14}$ ,  $NR^{13}SO_2R^{14}$ ,  $NR^{13}SONR^{14}R^{15}$ ,  $NR^{13}SO_2NR^{14}R^{15}$ ,  $P(O)R^{13}R^{14}$ ,  $P^+R^{13}R^{14}R^{15}A^-$ ,  $P(OR^{13})OR^{14}$ ,  $S^+R^{13}R^{14}A^-$ , and  $N^+R^9R^{11}R^{12}A^-$ ,

wherein:

$A^-$  is a pharmaceutically acceptable anion and  $M$  is a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of  $OR^7$ ,  $NR^7R^8$ ,  $SR^7$ ,  $S(O)R^7$ ,  $SO_2R^7$ ,  $SO_3R^7$ ,  $CO_2R^7$ ,  $CN$ , oxo,  $CONR^7R^8$ ,  $N^+R^7R^8R^9A^-$ , alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl,  $P(O)R^7R^8$ ,  $P^+R^7R^8R^9A^-$ , and  $P(O)(OR^7)OR^8$ , and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,  $NR^7$ ,  $N^+R^7R^8A^-$ ,  $S$ ,  $SO$ ,  $SO_2$ ,  $S^+R^7A^-$ ,  $PR^7$ ,  $P(O)R^7$ ,

$P^+R^7R^8A^-$ , or phenylene, and  $R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, arylalkyl, alkylarylalkyl, alkylheteroarylalkyl, alkylheterocyclalkyl, cycloalkyl, heterocycle, heteroaryl, quaternary heterocycle, quaternary heteroaryl, heterocyclalkyl, heteroarylalkyl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, alkylammoniumalkyl, and carboxyalkylaminocarbonylalkyl,

wherein alkyl, alkenyl, alkynyl, arylalkyl, heterocycle, and polyalkyl optionally have one or more carbons replaced by O,  $NR^9$ ,  $N^+R^9R^{10}A^-$ , S, SO, SO<sub>2</sub>,  $S^+R^9A^-$ ,  $PR^9$ ,  $P^+R^9R^{10}A^-$ ,  $P(O)R^9$ , phenylene, carbohydrate, amino acid, peptide, or polypeptide, and

$R^{13}$ ,  $R^{14}$ , and  $R^{15}$  are optionally substituted with one or more groups selected from the group consisting of hydroxy, amino, sulfo, carboxy, alkyl, carboxyalkyl, heterocycle, heteroaryl, sulfoalkyl, quaternary heterocycle, quaternary heteroaryl, quaternary heterocyclalkyl, quaternary heteroarylalkyl, guanidinyl,  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ , CN, halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{10}R^{11}A^-$ ,  $S^+R^9R^{10}A^-$ , and  $C(O)OM$ ,

wherein  $R^{16}$  and  $R^{17}$  are independently selected from the substituents constituting  $R^9$  and M; or

$R^{13}$  and  $R^{14}$ , together with the nitrogen atom to which they are attached form a mono- or polycyclic heterocycle that is optionally substituted with one or more radicals selected from the group consisting of oxo, carboxy and quaternary salts; or

$R^{14}$  and  $R^{15}$ , together with the nitrogen atom to which they are attached, form a cyclic ring; and

$R^{30}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, arylalkyl, carboxyalkyl, carboxyheteroaryl, carboxyheterocycle, carboalkoxyalkyl, carboxyalkylamino, heteroarylalkyl, heterocyclalkyl, and alkylammoniumalkyl; and

$R^7$  and  $R^8$  are hydrogen; and

one or more  $R^x$  are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heteroaryl, polyether, quaternary heterocycle, quaternary heteroaryl,  $OR^{13}$ ,  $NR^{13}R^{14}$ ,  $SR^{13}$ ,  $S(O)R^{13}$ ,  $S(O)_2R^{13}$ ,  $SO_3R^{13}$ ,  $S^+R^{13}R^{14}A^-$ ,  $NR^{13}OR^{14}$ ,  $NR^{13}NR^{14}R^{15}$ ,  $NO_2$ ,  $CO_2R^{13}$ ,  $CN$ ,  $OM$ ,  $SO_2OM$ ,  $SO_2NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)NR^{13}R^{14}$ ,  $NR^{14}C(O)R^{13}$ ,  $C(O)OM$ ,  $COR^{13}$ ,  $OR^{18}$ ,  $S(O)_nNR^{18}$ ,  $NR^{13}R^{18}$ ,  $NR^{18}OR^{14}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $P^+R^9R^{11}R^{12}A^-$ , amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,  $SO_3R^9$ , oxo,  $CO_2R^9$ ,  $CN$ , halogen,  $CONR^9R^{10}$ ,  $SO_2OM$ ,  $SO_2NR^9R^{10}$ ,  $PO(OR^{16})OR^{17}$ ,  $P^+R^9R^{11}R^{12}A^-$ ,  $S^+R^9R^{10}A^-$ , or  $C(O)OM$ , and

wherein  $R^{18}$  is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, and alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heteroaryl, alkyl, quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituents selected from the group consisting of  $OR^9$ ,  $NR^9R^{10}$ ,  $N^+R^9R^{11}R^{12}A^-$ ,  $SR^9$ ,  $S(O)R^9$ ,  $SO_2R^9$ ,

SRL 6071  
(C-3021/2)

663

$\text{SO}_3\text{R}^9$ , oxo,  $\text{CO}_2\text{R}^9$ , CN, halogen,  $\text{CONR}^9\text{R}^{10}$ ,  $\text{SO}_3\text{R}^9$ ,  $\text{SO}_2\text{OM}$ ,  
 $\text{SO}_2\text{NR}^9\text{R}^{10}$ ,  $\text{PO}(\text{OR}^{16})\text{OR}^{17}$ , and  $\text{C}(\text{O})\text{OM}$ ,


wherein in  $\text{R}^x$ , one or more carbons are optionally  
replaced by O,  $\text{NR}^{13}$ ,  $\text{N}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , S, SO,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^{13}\text{A}^-$ ,  $\text{PR}^{13}$ ,  
5  $\text{P}(\text{O})\text{R}^{13}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , phenylene, amino acid, peptide,  
polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid,  
peptide, polypeptide, and carbohydrate, one or more  
carbons are optionally replaced by O,  $\text{NR}^9$ ,  $\text{N}^+\text{R}^9\text{R}^{10}\text{A}^-$ , S,  
10  $\text{SO}$ ,  $\text{SO}_2$ ,  $\text{S}^+\text{R}^9\text{A}^-$ ,  $\text{PR}^9$ ,  $\text{P}^+\text{R}^9\text{R}^{10}\text{A}^-$ , or  $\text{P}(\text{O})\text{R}^9$ ;

wherein quaternary heterocycle and quaternary  
heteroaryl are optionally substituted with one or more  
groups selected from the group consisting of alkyl,  
alkenyl, alkynyl, polyalkyl, polyether, aryl,  
15 haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,  
oxo,  $\text{OR}^{13}$ ,  $\text{NR}^{13}\text{R}^{14}$ ,  $\text{SR}^{13}$ ,  $\text{S}(\text{O})\text{R}^{13}$ ,  $\text{SO}_2\text{R}^{13}$ ,  $\text{SO}_3\text{R}^{13}$ ,  $\text{NR}^{13}\text{OR}^{14}$ ,  
 $\text{NR}^{13}\text{NR}^{14}\text{R}^{15}$ ,  $\text{NO}_2$ ,  $\text{CO}_2\text{R}^{13}$ , CN, OM,  $\text{SO}_2\text{OM}$ ,  $\text{SO}_2\text{NR}^{13}\text{R}^{14}$ ,  
 $\text{C}(\text{O})\text{NR}^{13}\text{R}^{14}$ ,  $\text{C}(\text{O})\text{OM}$ ,  $\text{COR}^{13}$ ,  $\text{P}(\text{O})\text{R}^{13}\text{R}^{14}$ ,  $\text{P}^+\text{R}^{13}\text{R}^{14}\text{R}^{15}\text{A}^-$ ,  
20  $\text{P}(\text{OR}^{13})\text{OR}^{14}$ ,  $\text{S}^+\text{R}^{13}\text{R}^{14}\text{A}^-$ , and  $\text{N}^+\text{R}^9\text{R}^{11}\text{R}^{12}\text{A}^-$ ; and

$\text{R}^e$  is an electron-withdrawing group located at the  
para or ortho position.

336. The process of claim 335 wherein the cyclic  
sulfate has the formula:




XL

and the halobenzene has the formula:

SRL 6071  
(C-3021/2)

664



wherein  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^x$  and  $R^e$  are as defined in claim 335, and  $R^h$  is halogen.

5            337. The process of claim 336 wherein the sulfate group is removed by treating the intermediate with a hydrolyzing agent.

10           338. The process of claim 337 wherein the hydrolyzing agent is a mineral acid.

15           339. The process of claim 336 wherein the hydrolyzing agent is selected from the group consisting of hydrochloric acid and sulfuric acid.

            340. The process of claim 336 wherein the abstracting agent is a dialkali metal sulfide.

20           341. The process of claim 336 wherein the abstracting agent is dilithium sulfide.

            342. The process of claim 336 wherein  $R^1$  and  $R^2$  are alkyl.

25           343. The process of claim 336 wherein  $R^1$  and  $R^2$  are selected from the group consisting of ethyl, n-butyl, iso-butyl and pentyl.

30           344. The process of claim 336 wherein  $R^1$  and  $R^2$  are n-butyl.

345. The process of claim 336 wherein  $R^h$  is chloro.

5 346. The process of claim 336 wherein  $R^e$  is p-nitro.

10 347. The process of claim 336 wherein the alcohol is oxidized with an oxidizing agent to form a sulfone.

15 348. The process of claim 336 wherein the sulfone is oxidized with an oxidizing agent to form a sulfone-aldehyde.

20 349. The process of claim 336 wherein the sulfone-aldehyde is cyclized with a cyclizing agent that is a base having a pH between about 8 to about 9.

25 350. The process of claim 336 wherein the sulfone-aldehyde is cyclized with a cyclizing agent that is an alkali alkoxide base.

30 351. The process of claim 336 wherein the sulfone-aldehyde is cyclized with potassium tert-butoxide.

35 352. The process of claim 336 wherein the alcohol is oxidized with metachloroperbenzoic acid to form a sulfone; the aldehyde is oxidized with pyridinium chlorochromate to form a sulfone-aldehyde; and the sulfone-aldehyde is cyclized with potassium tert-butoxide.